

DEFORMATIONS OF GRADED NILPOTENT LIE ALGEBRAS AND SYMPLECTIC STRUCTURES

DMITRI V. MILLIONSCHIKOV

ABSTRACT. We study symplectic structures on filiform Lie algebras – nilpotent Lie algebras of the maximal length of the descending central sequence. There are two basic examples of symplectic $\mathbb{Z}_{>0}$ -graded filiform Lie algebras defined by their bases $e_1, \dots, e_{2k}$ and structure relations

- 1) $\mathfrak{m}_0(2k) : [e_1, e_i] = e_{i+1}, i = 2, \dots, 2k-1.$
- 2) $\mathcal{V}_{2k} : [e_i, e_j] = (j-i)e_{i+j}, i+j \leq 2k.$

Let $\mathfrak{g}$ be a symplectic filiform Lie algebra and $\dim \mathfrak{g} = 2k \geq 12$. Then $\mathfrak{g}$ is isomorphic to some $\mathbb{Z}_{>0}$ -filtered deformation either of $\mathfrak{m}_0(2k)$ or of $\mathcal{V}_{2k}$. In the present article we classify $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$, i.e., Lie algebras with structure relations of the following form:

$$[e_i, e_j] = (j-i)e_{i+j} + \sum_{l=1}^l c_{ij}^l e_{i+j+l}, \quad i+j \leq n$$

Namely we prove that for $n \geq 16$ the moduli space $\mathcal{M}_n$ of these algebras can be identified with the orbit space of the following $\mathbb{K}^*$ -action on $\mathbb{K}^5$:

$$\alpha \star X = (\alpha^{n-11} x_1, \alpha^{n-10} x_2, \dots, \alpha^{n-7} x_5), \alpha \in \mathbb{K}^*, X \in \mathbb{K}^5.$$

For $n = 2k$ the subspace $\mathcal{M}_{2k}^{symp} \subset \mathcal{M}_{2k}$ of symplectic Lie algebras is determined by equation $x_1 = 0$. A table with the structure constants of symplecto-isomorphism classes in $\mathcal{M}_{2k}^{symp}$ is presented.

INTRODUCTION

Nilmanifolds $M = G/\Gamma$ (compact homogeneous spaces of nilpotent Lie groups G over lattices Γ) are important examples of symplectic manifolds that do not admit Kähler structures. An interesting family of graded symplectic nilpotent Lie algebras $\mathcal{V}_{2k}$ (and corresponding family of nilmanifolds M_n) was considered in [1], [2], [3]. The finite dimensional Lie algebras $\mathcal{V}_n$, that are defined by the commuting relations $[e_i, e_j] = (j-i)e_{i+j}, i+j \leq n$, "came" from the infinite dimensional Virasoro algebra and they are examples of so-called filiform Lie algebras – nilpotent Lie algebras $\mathfrak{g}$ with the maximal length $s = \dim \mathfrak{g} - 1$ of the descending central sequence of $\mathfrak{g}$. The study of filiform Lie algebras was started by M. Vergne in [22], [23].

The classification of symplectic filiform Lie algebras of dimensions ≤ 10 was discussed in [10], [11]. The present paper is the continuation of [19], where a criterion of the existence of a symplectic structure on a filiform Lie algebra $\mathfrak{g}$ was proposed. In particular in a symplectic filiform $\mathfrak{g} = L^1 \mathfrak{g}$ one can find the ideal $L^2 \mathfrak{g}$ of codimension 1 such that the sequence of ideals $L^i \mathfrak{g}, i = 1, \dots, 2k$, where $L^i \mathfrak{g} = C^{i-1} \mathfrak{g}, i = 3, \dots, n$ ($\{C^i \mathfrak{g}\}$ are the ideals of the descending central sequence C of $\mathfrak{g}$) determines a decreasing filtration L of the Lie algebra $\mathfrak{g}$. The associated

1991 *Mathematics Subject Classification.* 17B30, 17B56, 17B70, 53D.

Partially supported by the Russian Foundation for Fundamental Research, grant no. 99-01-00090 and PAI-RUSSIE, dossier no. 04495UL.

graded Lie algebra $\text{gr}_L \mathfrak{g}$ is symplectic also. The graded filiform algebras of the type $\text{gr}_L \mathfrak{g}$ were classified in [18], [19]. There are two one-parameter families of graded symplectic filiform Lie algebras of dimensions 8, 10. But if $\text{gr}_L \mathfrak{g}$ is a symplectic filiform Lie algebra with $\dim \mathfrak{g} = 2k \geq 12$ then $\text{gr}_L \mathfrak{g}$ is isomorphic either to $\mathfrak{m}_0(2k)$ or to $\mathcal{V}_{2k}$. In other words in dimensions $2k \geq 12$ one can obtain symplectic filiform Lie algebras as special deformations (that we call $\mathbb{Z}_{>0}$ -filtered deformations) of two graded Lie algebras: $\mathfrak{m}_0(2k)$ and $\mathcal{V}_{2k}$.

We classify $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$, i.e., Lie algebras with the structure relations of the following form:

$$[e_i, e_j] = (j-i)e_{i+j} + \sum_{l=1}^j c_{ij}^l e_{i+j+l}, \quad i+j \leq n.$$

We compute in the Section 6 the space $H^2(\mathcal{V}_n, \mathcal{V}_n)$ for $n \geq 12$. To the $\mathbb{Z}_{>0}$ -filtered deformations corresponds the subspace $\oplus_{>0} H_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$. The main theorem 7.9 of the present article asserts that for $n \geq 16$ the moduli space $\mathcal{M}_n$ (i.e. the set of isomorphism classes) of these algebras can be identified with the orbit space of the following $\mathbb{K}^*$ -action on $\mathbb{K}^5 = \oplus_{>0} H_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$:

$$\alpha \star X = (\alpha^{n-11}x_1, \alpha^{n-10}x_2, \dots, \alpha^{n-7}x_5), \alpha \in \mathbb{K}^*, X \in \mathbb{K}^5,$$

where the coordinates $x_1, \dots, x_5$ of the space $\mathbb{K}^5 = \oplus_{>0} H_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$ are defined by the choice of the basic cocycles $\psi_{n,12-i}, i = 1, \dots, 5$.

For $n = 2k$ the subspace $\mathcal{M}_{2k}^{\text{sympl}} \subset \mathcal{M}_{2k}$ of symplectic Lie algebras is determined by equation $x_1 = 0$. A table with structure constants of symplecto-isomorphism classes in $\mathcal{M}_{2k}^{\text{sympl}}$ is presented in the Section 9.

1. INVARIANT SYMPLECTIC STRUCTURES ON LIE GROUPS

Definition 1.1. A Lie group G is said to have a left-invariant symplectic structure if it has a left-invariant non-degenerate closed 2-form ω .

Example 1.2. G is a two-dimensional abelian Lie group $\mathbb{R}^2$ with coordinates x, y and $\omega = dx \wedge dy$.

Example 1.3. G is a direct product $\mathcal{H}_3 \times \mathbb{R}$ of the Heisenberg group $\mathcal{H}_3$ of all matrices of the form

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}, \quad x, y, z \in \mathbb{R},$$

and abelian $\mathbb{R}$ (with coordinate t). The invariant symplectic form ω is defined as

$$\omega = dx \wedge (dz - xdy) + dy \wedge dt.$$

Theorem 1.4 (B.-Y. Chu, [4]). *Any semisimple Lie group has no left-invariant symplectic structure.*

Theorem 1.5 (B.-Y. Chu, [4]). *A connected unimodular Lie group admitting a left-invariant symplectic structure must to be solvable.*

One can obtain examples of left-invariant symplectic structures in the framework of Kirillov's orbit method.

Let us consider the coadjoint action Ad^* of a Lie group G on the dual $\mathfrak{g}^*$ of its Lie algebra $\mathfrak{g}$:

$$(Ad^*(g)f)(X) = f(Ad(g^{-1})X), \quad \forall X \in \mathfrak{g},$$

where $g \in G, f \in \mathfrak{g}^*$. The orbit O_f of this action has a homogeneous symplectic structure ω_{O_f} such that

$$\pi_f^*(\omega_{O_f}) = df,$$

where π_f denotes the natural mapping $\pi_f : G \rightarrow O_f, \pi_f(g) = Ad^*(g)f$. Let the stabilizer $G_f = \{g \in G | Ad^*(g)f = f\}$ of $f \in \mathfrak{g}^*$ be a normal subgroup of G then the orbit O_f can be identified with the quotient group G/G_f and the corresponding symplectic structure ω_{O_f} is left G/G_f -invariant. If G is a nilpotent Lie group then nilpotent ones are G_f and G/G_f .

Definition 1.6. A skew-symmetric non-degenerate bilinear form ω on the Lie algebra $\mathfrak{g}$ is called symplectic if it closed, i.e.

$$\omega([x, y], z) + \omega([y, z], x) + \omega([z, x], y) = 0, \quad \forall x, y, z \in \mathfrak{g}.$$

If ω_G is a left-invariant symplectic form on G , then ω_G defines a symplectic structure $\omega_{\mathfrak{g}}$ on the Lie algebra $\mathfrak{g}$ of G , and conversely any symplectic form $\omega_{\mathfrak{g}}$ of $\mathfrak{g}$ defines a left-invariant symplectic structure on G .

Lemma 1.7 (A. Médina, P. Revoy [15]). *Let $\mathfrak{g}$ be a symplectic Lie algebra with non-trivial center $Z(\mathfrak{g})$ and I be a one-dimensional subspace in $Z(\mathfrak{g})$ and I^ω its symplectic complement with respect to ω . Then one can consider two following exact sequences of Lie algebras and their homomorphisms*

$$(1) \quad \begin{aligned} &0 \rightarrow I \rightarrow I^\omega \rightarrow I^\omega/I \rightarrow 0 \\ &0 \rightarrow I^\omega \rightarrow \mathfrak{g} \rightarrow \mathfrak{g}/I^\omega \cong I \rightarrow 0 \end{aligned}$$

where I^ω/I is symplectic Lie algebra (the restriction of ω to I^ω defines symplectic form $\tilde{\omega}$ on the quotient-algebra I^ω/I).

In other words $2k$ -dimensional symplectic Lie algebra $\mathfrak{g}$ with non-trivial center can be obtained from $2k-2$ -dimensional symplectic I^ω/I by means of two consecutive operations:

- 1) one-dimensional central extension of I^ω/I by I ;
- 2) semidirect product of I^ω and one-dimensional $\mathfrak{g}/I^\omega \cong I$.

The combination of these two operations is called the double extension of symplectic Lie algebra I^ω/I .

Theorem 1.8 (A. Médina, P. Revoy [15]). *Symplectic nilpotent Lie algebras can be obtained by means of the sequence of consecutive double extensions starting with trivial Lie algebra of zero dimension.*

The even-dimensional nilpotent Lie algebras are classified for dimensions $2k \leq 6$ ([20]). The classification of symplectic 6-dimensional nilpotent Lie algebras based on this classification was done in [11].

2. FILIFORM LIE ALGEBRAS

The sequence of ideals of a Lie algebra $\mathfrak{g}$

$$C^1 \mathfrak{g} = \mathfrak{g} \supset C^2 \mathfrak{g} = [\mathfrak{g}, \mathfrak{g}] \supset \dots \supset C^k \mathfrak{g} = [\mathfrak{g}, C^{k-1} \mathfrak{g}] \supset \dots$$

is called the descending central sequence of $\mathfrak{g}$.

A Lie algebra $\mathfrak{g}$ is called nilpotent if there exists s such that:

$$C^{s+1} \mathfrak{g} = [\mathfrak{g}, C^s \mathfrak{g}] = 0, \quad C^s \mathfrak{g} \neq 0.$$

The natural number s is called the nil-index of the nilpotent Lie algebra $\mathfrak{g}$, or $\mathfrak{g}$ is called s -step nilpotent Lie algebra.

Let $\mathfrak{g}$ be a Lie algebra. We call a set F of subspaces

$$\mathfrak{g} \supset \cdots \supset F^i \mathfrak{g} \supset F^{i+1} \mathfrak{g} \supset \cdots \quad (i \in \mathbb{Z})$$

a decreasing filtration F of $\mathfrak{g}$ if F is compatible with the Lie structure

$$[F^k \mathfrak{g}, F^l \mathfrak{g}] \subset F^{k+l} \mathfrak{g}, \quad \forall k, l \in \mathbb{Z}.$$

Let $\mathfrak{g}$ be a filtered Lie algebra. A graded Lie algebra

$$\mathrm{gr}_F \mathfrak{g} = \bigoplus_{k=1} (\mathrm{gr}_F \mathfrak{g})_k, \quad (\mathrm{gr}_F \mathfrak{g})_k = F^k \mathfrak{g} / F^{k+1} \mathfrak{g}$$

is called the associated graded Lie algebra $\mathrm{gr}_F \mathfrak{g}$.

The ideals $C^k \mathfrak{g}$ of the descending central sequence define a decreasing filtration C of the Lie algebra $\mathfrak{g}$

$$C^1 \mathfrak{g} = \mathfrak{g} \supset C^2 \mathfrak{g} \supset \cdots \supset C^k \mathfrak{g} \supset \cdots; \quad [C^k \mathfrak{g}, C^l \mathfrak{g}] \subset C^{k+l} \mathfrak{g}.$$

One can consider the associated graded Lie algebra $\mathrm{gr}_C \mathfrak{g}$.

The finite filtration C of a nilpotent Lie algebra $\mathfrak{g}$ is called the canonical filtration of a nilpotent Lie algebra $\mathfrak{g}$.

Proposition 2.1. *Let $\mathfrak{g}$ be a n -dimensional nilpotent Lie algebra. Then for its nil-index we have the estimate $s \leq n - 1$.*

Definition 2.2. A nilpotent n -dimensional Lie algebra $\mathfrak{g}$ is called filiform Lie algebra if it has the nil-index $s = n - 1$.

Example 2.3. The Lie algebra $\mathfrak{m}_0(n)$ is defined by its basis $e_1, e_2, \dots, e_n$ with commuting relations:

$$[e_1, e_i] = e_{i+1}, \quad \forall 2 \leq i \leq n-1.$$

Remark. We will omit in the sequel trivial commuting relations $[e_i, e_j] = 0$ in the definitions of Lie algebras.

Example 2.4. The Lie algebra $\mathfrak{m}_2(n)$ is defined by its basis $e_1, e_2, \dots, e_n$ and commuting relations:

$$[e_1, e_i] = e_{i+1}, \quad 2 \leq i \leq n-1; \quad [e_2, e_j] = e_{j+2}, \quad 3 \leq j \leq n-2.$$

Example 2.5. Let us define the algebra L_k as the infinite-dimensional Lie algebra of polynomial vector fields on the real line $\mathbb{R}^1$ with a zero in $x = 0$ of order not less than $k + 1$.

The algebra L_k can be defined by its infinite basis and commuting relations

$$e_i = x^{i+1} \frac{d}{dx}, \quad i \in \mathbb{N}, \quad i \geq k, \quad [e_i, e_j] = (j - i)e_{i+j}, \quad \forall i, j \in \mathbb{N}.$$

One can consider the n -dimensional quotient algebra $\mathcal{V}_n = L_1 / L_{n+1}$.

The Lie algebras $\mathfrak{m}_0(n)$, $\mathfrak{m}_2(n)$, $\mathcal{V}_n$ considered above are filiform Lie algebras.

Proposition 2.6. *Let $\mathfrak{g}$ be a filiform Lie algebra and $\mathrm{gr}_C \mathfrak{g} = \bigoplus_i (\mathrm{gr}_C \mathfrak{g})_i$ is the corresponding associated (with respect to the canonical filtration C) graded Lie algebra. Then*

$$\dim(\mathrm{gr}_C \mathfrak{g})_1 = 2, \quad \dim(\mathrm{gr}_C \mathfrak{g})_2 = \cdots = \dim(\mathrm{gr}_C \mathfrak{g})_{n-1} = 1.$$

We have the following isomorphisms of graded Lie algebras:

$$\mathrm{gr}_C \mathfrak{m}_2(n) \cong \mathrm{gr}_C \mathcal{V}_n \cong \mathrm{gr}_C \mathfrak{m}_0(n) \cong \mathfrak{m}_0(n).$$

Theorem 2.7 (M. Vergne [23]). *Let $\mathfrak{g} = \oplus_{\alpha} \mathfrak{g}_{\alpha}$ be a graded n -dimensional filiform Lie algebra and*

$$(2) \quad \dim \mathfrak{g}_1 = 2, \quad \dim \mathfrak{g}_2 = \cdots = \dim \mathfrak{g}_{n-1} = 1.$$

then

- 1) if $n = 2k + 1$, then $\mathfrak{g}$ is isomorphic to $\mathfrak{m}_0(2k + 1)$;
- 2) if $n = 2k$, then $\mathfrak{g}$ is isomorphic either to $\mathfrak{m}_0(2k)$ or to the Lie algebra $\mathfrak{m}_1(2k)$, defined by its basis $e_1, \dots, e_{2k}$ and commutating relations:

$$[e_1, e_i] = e_{i+1}, \quad i = 2, \dots, 2k-1; \quad [e_j, e_{2k+1-j}] = (-1)^{j+1} e_{2k}, \quad j = 2, \dots, k.$$

Remark. In the settings of the Theorem 2.7 the gradings of the algebras $\mathfrak{m}_0(n)$, $\mathfrak{m}_1(n)$ are defined as $\mathfrak{g}_1 = \mathrm{Span}(e_1, e_2)$, $\mathfrak{g}_i = \mathrm{Span}(e_{i+1})$, $i = 2, \dots, n-1$.

Corollary 2.8 (M. Vergne [23]). *Let $\mathfrak{g}$ be a filiform Lie algebra. Then one can choose a so-called adapted basis $e_1, e_2, \dots, e_n$ in $\mathfrak{g}$:*

$$(3) \quad [e_1, e_i] = e_{i+1}, \quad i=2, \dots, n-1; \quad [e_i, e_j] = \begin{cases} \sum_{k=0}^{n-i-j} c_{ij}^{i+j+k} e_{i+j+k}, & i+j \leq n; \\ (-1)^i \alpha e_n, & i+j = n+1; \\ 0, & i+j > n+1; \\ & 2 \leq i < j \leq n. \end{cases}$$

where $\alpha = 0$ if n is odd number.

3. SYMPLECTIC FILIFORM LIE ALGEBRAS: FILTRATIONS AND GRADINGS

Definition 3.1. A Lie algebra $\mathfrak{g}$ is called symplectic if it admits at least one symplectic structure.

Lemma 3.2 ([19]). *Let $\mathfrak{g}$ be an $2k$ -dimensional symplectic filiform Lie algebra, then*

$$(4) \quad \mathrm{gr}_C \mathfrak{g} \cong \mathfrak{m}_0(2k).$$

In other words: let $\mathfrak{g}$ be a symplectic filiform Lie algebra and $e_1, e_2, \dots, e_{2k}$ be some adapted basis, then $[e_i, e_j] = 0$, $i+j = 2k+1$.

Proposition 3.3. *A fixed adapted basis of $\mathfrak{g}$ $e_1, e_2, \dots, e_{2k}$ such that $[e_i, e_j] = 0$, $i+j = 2k+1$ defines a non-canonical filtration L of $\mathfrak{g}$:*

$$(5) \quad \begin{aligned} \mathfrak{g} &= L^1 \mathfrak{g} \supset L^2 \mathfrak{g} \supset \cdots \supset L^{2k} \mathfrak{g} \supset \{0\}, \\ L^j \mathfrak{g} &= \mathrm{Span}(e_j, \dots, e_{2k}), \quad j = 1, \dots, 2k. \end{aligned}$$

Remark. For homogeneous components of associated graded $\mathrm{gr}_L \mathfrak{g}$ we have

$$\dim(\mathrm{gr}_L \mathfrak{g})_1 = \dim(\mathrm{gr}_L \mathfrak{g})_2 = \cdots = \dim(\mathrm{gr}_L \mathfrak{g})_{2k} = 1.$$

Proposition 3.4 ([19]). *Let $\mathfrak{g}$ be a symplectic filiform Lie algebra with a fixed adapted basis. Then the corresponding associated graded Lie algebra $\mathrm{gr}_L \mathfrak{g} = \oplus_i (\mathrm{gr}_L \mathfrak{g})_i$ is symplectic also.*

Remark. It was shown in [19] that the previous condition is necessary but not sufficient condition.

Theorem 3.5 (D. Millionschikov, [19]). *Let $\mathfrak{g} = \bigoplus_{\alpha=1}^{2k} \mathfrak{g}_\alpha$ be a real graded symplectic Lie algebra such that*

$$(6) \quad \dim \mathfrak{g}_\alpha = 1, \quad 1 \leq \alpha \leq 2k; \quad [\mathfrak{g}_1, \mathfrak{g}_\alpha] = \mathfrak{g}_{\alpha+1}, \quad 2 \leq \alpha \leq 2k-1.$$

Then $\mathfrak{g}$ is isomorphic to the one and only one Lie algebra from the following list :

<i>dim</i>	<i>algebra</i>	<i>commutating relations</i>	<i>symplectic form</i>
4	$\mathfrak{m}_0(4)$	$[e_1, e_2] = e_3, [e_1, e_3] = e_4$	$e^1 \wedge e^4 - e^2 \wedge e^3$
6	$\mathfrak{m}_0(6)$	$[e_1, e_i] = e_{i+1}, i = 2, \dots, 5$	$e^1 \wedge e^6 - e^2 \wedge e^5 + e^3 \wedge e^4$
	$\mathcal{V}_6$	$[e_i, e_j] = (j-i)e_{i+j}, i+j \leq 6$	$5e^1 \wedge e^6 + 3e^2 \wedge e^5 + e^3 \wedge e^4$
8	$\mathfrak{m}_0(8)$	$[e_1, e_i] = e_{i+1}, i = 2, \dots, 7$	$e^1 \wedge e^8 - e^2 \wedge e^7 + e^3 \wedge e^6 - e^4 \wedge e^5$
	$\mathfrak{g}_{8,\alpha}$	(1) $\alpha \neq -\frac{5}{2}, -2, -\frac{1}{2}, \frac{1}{2}$	$e^1 \wedge e^8 + \frac{2\alpha^2+3\alpha-2}{2\alpha+5} e^2 \wedge e^7 +$ $+ \frac{2\alpha+2}{2\alpha+5} e^3 \wedge e^6 + \frac{3}{2\alpha+5} e^4 \wedge e^5$
10	$\mathfrak{m}_0(10)$	$[e_1, e_i] = e_{i+1}, i = 2, \dots, 9$	$e^1 \wedge e^{10} - e^2 \wedge e^9 + e^3 \wedge e^8 - e^4 \wedge e^7 + e^5 \wedge e^6$
	$\mathfrak{g}_{10,\alpha}$	$(1), (2)$ $\alpha \neq -\frac{5}{2}, -\frac{1}{4}, -1, -3, \alpha_1, \alpha_2;$ $\alpha_1, \alpha_2 \in \mathbb{R}, 2\alpha_1^3 + 2\alpha_1^2 + 3 = 0,$ $4\alpha_2^3 + 8\alpha_2^2 - 8\alpha_2 - 21 = 0$	$e^1 \wedge e^{10} + \frac{2\alpha^3+2\alpha^2+3}{2(\alpha^2+4\alpha+3)} e^2 \wedge e^9 +$ $+ \frac{4\alpha^3+8\alpha^2-8\alpha-21}{2(\alpha^2+4\alpha+3)(2\alpha+5)} e^3 \wedge e^8 +$ $+ \frac{3(2\alpha^2+4\alpha+5)}{2(\alpha^2+4\alpha+3)(2\alpha+5)} e^4 \wedge e^7 +$ $+ \frac{3(4\alpha+1)}{2(\alpha^2+4\alpha+3)(2\alpha+5)} e^5 \wedge e^6$
$2k \geq 12$	$\mathfrak{m}_0(2k)$	$[e_1, e_i] = e_{i+1}, i = 2, \dots, 2k-1$	$\frac{1}{2} \sum_{i+j=2k+1} (-1)^{i+1} e^i \wedge e^j$
	$\mathcal{V}_{2k}$	$[e_i, e_j] = (j-i)e_{i+j}, i+j \leq 2k$	$\frac{1}{2} \sum_{i+j=2k+1} (j-i) e^i \wedge e^j$

	<i>commutating relations for $\mathfrak{g}_{8,\alpha}$ and $\mathfrak{g}_{10,\alpha}$</i>
(1)	$[e_1, e_2] = e_3, [e_1, e_3] = e_4, [e_1, e_4] = e_5, [e_1, e_5] = e_6, [e_1, e_6] = e_7,$ $[e_2, e_3] = (2+\alpha)e_5, [e_2, e_4] = (2+\alpha)e_6, [e_2, e_5] = (1+\alpha)e_7, [e_3, e_4] = e_7$ $[e_1, e_7] = e_8, [e_2, e_6] = \alpha e_8, [e_3, e_5] = e_8$
(2)	$[e_1, e_8] = e_9, [e_2, e_7] = \frac{2\alpha^2+3\alpha-2}{2\alpha+5} e_9, [e_3, e_6] = \frac{2\alpha+2}{2\alpha+5} e_9, [e_4, e_5] = \frac{3}{2\alpha+5} e_9$ $[e_1, e_9] = e_{10}, [e_2, e_8] = \frac{2\alpha^2+\alpha-1}{2\alpha+5} e_{10}, [e_3, e_7] = \frac{2\alpha-1}{2\alpha+5} e_{10}, [e_4, e_6] = \frac{3}{2\alpha+5} e_{10}$

Remark. In the fourth column of the table we give only one variant of possible symplectic structure.

Corollary 3.6. *Let $\mathfrak{g}$ be a symplectic filiform Lie algebra of dimension $2k \geq 12$ then one can choose a basis $e_1, \dots, e_{2k}$ in $\mathfrak{g}$ such that the corresponding commutating relations will be either*

$$(7) \quad [e_1, e_i] = e_{i+1} + \sum_{l=1}^{2k-i-1} c_{1i}^l e_{i+l+1}, \quad i = 2, \dots, 2k-1;$$

$$[e_i, e_j] = \sum_{l=1}^{2k-i-j} c_{ij}^l e_{i+j+l}, \quad i+j \leq 2k; 2 \leq i < j \leq 2k;$$

or

$$(8) \quad [e_i, e_j] = (j-i)e_{i+j} + \sum_{l=1}^{2k-i-j} c_{ij}^l e_{i+j+l}, \quad i+j \leq 2k; 1 \leq i < j \leq 2k;$$

Remark. The one-parameter family $\mathfrak{g}_{8,\alpha}$ was considered in [11] as well as corresponding symplectic form $\omega_9(\alpha)$. In [10] symplectic (over $\mathbb{C}$) low-dimensional ($\dim \mathfrak{g} \leq 10$) filiform Lie algebras were classified (but this article contains some mistakes).

4. LIE ALGEBRA COHOMOLOGY

Let $\mathfrak{g}$ be a Lie algebra over $\mathbb{K}$ and $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ its linear representation (or in other words V is a $\mathfrak{g}$ -module). We denote by $C^q(\mathfrak{g}, V)$ the space of q -linear skew-symmetric mappings of $\mathfrak{g}$ into V . Then one can consider an algebraic complex:

$$V \xrightarrow{d_0} C^1(\mathfrak{g}, V) \xrightarrow{d_1} C^2(\mathfrak{g}, V) \xrightarrow{d_2} \dots \xrightarrow{d_{q-1}} C^q(\mathfrak{g}, V) \xrightarrow{d_q} \dots$$

where the differential d_q is defined by:

$$(9) \quad \begin{aligned} (d_q f)(X_1, \dots, X_{q+1}) &= \sum_{i=1}^{q+1} (-1)^{i+1} \rho(X_i)(f(X_1, \dots, \hat{X}_i, \dots, X_{q+1})) + \\ &+ \sum_{1 \leq i < j \leq q+1} (-1)^{i+j-1} f([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{q+1}). \end{aligned}$$

The cohomology of the complex $(C^*(\mathfrak{g}, V), d)$ is called the cohomology of the Lie algebra $\mathfrak{g}$ with coefficients in the representation $\rho : \mathfrak{g} \rightarrow V$.

In this article we will consider two main examples:

- 1) $V = \mathbb{K}$ and $\rho : \mathfrak{g} \rightarrow \mathbb{K}$ is trivial;
- 2) $V = \mathfrak{g}$ and $\rho = ad : \mathfrak{g} \rightarrow \mathfrak{g}$ is the adjoint representation of $\mathfrak{g}$.

The cohomology of $(C^*(\mathfrak{g}, \mathbb{K}), d)$ (the first example) is called the cohomology with trivial coefficients of the Lie algebra $\mathfrak{g}$ and is denoted by $H^*(\mathfrak{g})$. Also we fix the notation $H^*(\mathfrak{g}, \mathfrak{g})$ for the cohomology of $\mathfrak{g}$ with coefficients in the adjoint representation.

One can remark that $d_1 : C^1(\mathfrak{g}, \mathbb{K}) \rightarrow C^2(\mathfrak{g}, \mathbb{K})$ of the $(C^*(\mathfrak{g}, \mathbb{K}), d)$ is the dual mapping to the Lie bracket $[\cdot, \cdot] : \Lambda^2 \mathfrak{g} \rightarrow \mathfrak{g}$. Moreover the condition $d^2 = 0$ is equivalent to the Jacobi identity for $[\cdot, \cdot]$.

Let $\mathfrak{g} = \bigoplus_{\alpha} \mathfrak{g}_{\alpha}$ be a $\mathbb{Z}$ -graded Lie algebra and $V = \bigoplus_{\beta} V_{\beta}$ is a $\mathbb{Z}$ -graded $\mathfrak{g}$ -module, i.e., $\mathfrak{g}_{\alpha} V_{\beta} \subset V_{\alpha+\beta}$. Then the complex $(C^*(\mathfrak{g}, V), d)$ can be equipped with the $\mathbb{Z}$ -grading $C^q(\mathfrak{g}, V) = \bigoplus_{\mu} C_{(\mu)}^q(\mathfrak{g}, V)$, where a V -valued q -form c belongs to $C_{(\mu)}^q(\mathfrak{g}, V)$ iff for $X_1 \in \mathfrak{g}_{\alpha_1}, \dots, X_q \in \mathfrak{g}_{\alpha_q}$ we have

$$c(X_1, \dots, X_q) \in V_{\alpha_1 + \alpha_2 + \dots + \alpha_q + \mu}.$$

This grading is compatible with the differential d and hence we have $\mathbb{Z}$ -grading in cohomology:

$$H^q(\mathfrak{g}, V) = \bigoplus_{\mu \in \mathbb{Z}} H_{(\mu)}^q(\mathfrak{g}, V).$$

Remark. The trivial $\mathfrak{g}$ -module $\mathbb{K}$ has only one non-trivial homogeneous component $\mathbb{K} = \mathbb{K}_0$.

Example 4.1. Let $\mathfrak{g}$ be a Lie algebra with the basis $e_1, e_2, \dots, e_n$ and commuting relations

$$[e_i, e_j] = c_{ij} e_{i+j}, \quad i + j \leq n.$$

Let us consider the dual basis $e^1, e^2, \dots, e^n$. One can introduce a grading (that we will call the weight) of $\Lambda^*(\mathfrak{g}^*) = C^*(\mathfrak{g}, \mathbb{K})$:

$$\Lambda^*(\mathfrak{g}^*) = \bigoplus_{\lambda=1}^{n(n+1)/2} \Lambda_{(\lambda)}^*(\mathfrak{g}^*),$$

where a subspace $\Lambda_{(\lambda)}^q(\mathfrak{g}^*)$ is spanned by q -forms $\{e^{i_1} \wedge \dots \wedge e^{i_q}, i_1 + \dots + i_q = \lambda\}$. For instance a monomial $e^{i_1} \wedge \dots \wedge e^{i_q}$ has the degree q and the weight $\lambda = i_1 + \dots + i_q$.

The complex $(C^*(\mathfrak{g}, \mathfrak{g}), d)$ is $\mathbb{Z}$ -graded:

$$C^*(\mathfrak{g}, \mathfrak{g}) = \bigoplus_{\mu \in \mathbb{Z}} C_{(\mu)}^*(\mathfrak{g}, \mathfrak{g}),$$

where $C_{(\mu)}^q(\mathfrak{g}, \mathfrak{g})$ is spanned by monomials $\{e_l \otimes e^{i_1} \wedge \dots \wedge e^{i_q}, i_1 + \dots + i_q + \mu = l\}$.

Now we consider a filtered Lie algebra $\mathfrak{g}$ with a finite decreasing filtration F . One can define a decreasing filtration $\tilde{F}$ of $\Lambda^*(\mathfrak{g}^*)$.

$$\tilde{F}^\mu \Lambda^p(\mathfrak{g}^*) = \{\omega \in \Lambda^p(\mathfrak{g}^*) \mid \omega(F^{\alpha_1} \mathfrak{g} \wedge \dots \wedge F^{\alpha_p} \mathfrak{g}) = 0, \alpha_1 + \dots + \alpha_p + \mu \geq 0\}.$$

Example 4.2. Let $\mathfrak{g}$ be a Lie algebra with the basis $e_1, e_2, \dots, e_n$ and commuting relations

$$[e_i, e_j] = \sum_{k=0}^{n-i-j} c_{ij}^k e_{i+j+k}, i+j \leq n.$$

As it was remarked above the corresponding filtration L ($L^k = \text{Span}(e_k, \dots, e_n)$) of $\mathfrak{g}$ can be defined. The associated graded Lie algebra $\text{gr}_L \mathfrak{g}$ has the following structure relations

$$[e_i, e_j] = c_{ij}^0 e_{i+j}, i+j \leq n.$$

Let us consider the dual basis $e^1, e^2, \dots, e^n$. Then $\tilde{L}^\mu \Lambda^p(\mathfrak{g}^*)$ is spanned by p -monomials of weights less or equal to $-\mu$, i.e. by p -forms $e^{i_1} \wedge \dots \wedge e^{i_p}$ such that $i_1 + \dots + i_p \leq -\mu$. For instance

$$\tilde{L}^{-5} \Lambda^2(\mathfrak{g}^*) = \text{Span}(e^1 \wedge e^2, e^1 \wedge e^3, e^1 \wedge e^4, e^2 \wedge e^3).$$

Remark. One can consider the spectral sequence E_r that corresponds to the filtration $\tilde{F}$ of the complex $\Lambda^*(\mathfrak{g}^*)$. We have an isomorphism (see [23] for example)

$$E_1^{p,q} = H_{(-p)}^{p+q}(\text{gr}_F \mathfrak{g}).$$

Theorem 4.3 ([19]). *Let $\mathfrak{g}$ be a filiform Lie algebra such that $\text{gr}_C \mathfrak{g} \cong \mathfrak{m}_0(2k)$ and $\text{gr}_L \mathfrak{g}$ is symplectic.*

Then the Lie algebra $\mathfrak{g}$ is symplectic if and only if some homogeneous symplectic class $[\omega_{2k+1}] \in E_1^{-2k-1, 2k+3} = H_{(2k+1)}^2(\text{gr}_L \mathfrak{g})$ survives to the term E_∞ .

5. $\mathbb{Z}_{>0}$ -FILTERED DEFORMATIONS AND $H^2(\mathfrak{g}, \mathfrak{g})$

In this section we recall some definitions from the Nijenhuis-Richardson deformation theory (see [21]).

Definition 5.1. Let $\mathfrak{g}$ be a Lie algebra with a Lie bracket $[\cdot, \cdot]$ and $\Psi : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ is a skew-symmetric bilinear map. Ψ is called a deformation of $[\cdot, \cdot]$ iff $[\cdot, \cdot]' = [\cdot, \cdot] + \Psi$ is a Lie bracket on the vector space $\mathfrak{g}$.

The Jacobi identity for bracket $[\cdot]' = [\cdot] + \Psi$

$$[[x, y]', z]' + [[y, z]', x]' + [[z, x]', y]' = 0$$

is equivalent to the so-called deformation equation:

$$(10) \quad \begin{aligned} & \Psi([x, y], z) + \Psi([y, z], x) + \Psi([z, x], y) + [\Psi(x, y), z] + [\Psi(y, z), x] + [\Psi(y, z), x] + \\ & + \Psi(\Psi(x, y), z) + \Psi(\Psi(y, z), x) + \Psi(\Psi(z, x), y) = 0. \end{aligned}$$

The first six terms can be rewritten in the form $d\Psi(x, y, z)$ where $d : C^2(\mathfrak{g}, \mathfrak{g}) \rightarrow C^3(\mathfrak{g}, \mathfrak{g})$ is the differential of the complex $(C^*(\mathfrak{g}, \mathfrak{g}), d)$.

Finally we have

$$(11) \quad d\Psi + \frac{1}{2}[\Psi, \Psi] = 0,$$

where $[\cdot]$ denotes a symmetric bilinear function $[\cdot] : C^2(\mathfrak{g}, \mathfrak{g}) \times C^2(\mathfrak{g}, \mathfrak{g}) \rightarrow C^3(\mathfrak{g}, \mathfrak{g})$

$$(12) \quad \begin{aligned} [\Psi, \tilde{\Psi}](x, y, z) = & \Psi(\tilde{\Psi}(x, y), z) + \Psi(\tilde{\Psi}(y, z), x) + \Psi(\tilde{\Psi}(z, x), y) + \\ & + \tilde{\Psi}(\Psi(x, y), z) + \tilde{\Psi}(\Psi(y, z), x) + \tilde{\Psi}(\Psi(z, x), y). \end{aligned}$$

The last definition can be generalised in terms of Nijenhuis-Richardson bracket in $C^*(\mathfrak{g}, \mathfrak{g})$:

$$[\cdot] : C^p(\mathfrak{g}, \mathfrak{g}) \times C^q(\mathfrak{g}, \mathfrak{g}) \rightarrow C^{p+q-1}(\mathfrak{g}, \mathfrak{g}).$$

Namely, for $\alpha \in C^p(\mathfrak{g}, \mathfrak{g})$ and $\beta \in C^q(\mathfrak{g}, \mathfrak{g})$ one can define $[\alpha, \beta] \in C^{p+q-1}(\mathfrak{g}, \mathfrak{g})$:

$$(13) \quad \begin{aligned} [\alpha, \beta](\xi_1, \dots, \xi_{p+q-1}) = & \sum_{1 \leq i_1 < \dots < i_q \leq p+q-1} \alpha(\beta(\xi_{i_1}, \dots, \xi_{i_q}) \xi_1, \dots, \hat{\xi}_{i_1}, \dots, \hat{\xi}_{i_q}, \dots, \xi_{p+q-1}) + \\ & + (-1)^{pq+p+q} \sum_{1 \leq j_1 < \dots < j_p \leq p+q-1} \beta(\alpha(\xi_{j_1}, \dots, \xi_{j_p}) \xi_1, \dots, \hat{\xi}_{j_1}, \dots, \hat{\xi}_{j_p}, \dots, \xi_{p+q-1}). \end{aligned}$$

The Nijenhuis-Richardson bracket defines a Lie superalgebra structure in $C^*(\mathfrak{g}, \mathfrak{g})$, i.e., if $\alpha \in C^p(\mathfrak{g}, \mathfrak{g})$, $\beta \in C^q(\mathfrak{g}, \mathfrak{g})$ and $\gamma \in C^r(\mathfrak{g}, \mathfrak{g})$ then

$$(14) \quad \begin{aligned} 1) \quad & [\alpha, \beta] = -(-1)^{(p-1)(q-1)}[\beta, \alpha]; \\ 2) \quad & (-1)^{(p-1)(q-1)}[[\alpha, \beta], \gamma] + (-1)^{(q-1)(r-1)}[[\beta, \gamma], \alpha] + (-1)^{(r-1)(p-1)}[[\gamma, \alpha], \beta] = 0. \end{aligned}$$

Also we have the following important property:

$$d[\alpha, \beta] = [d\alpha, \beta] + (-1)^p[\alpha, d\beta].$$

Thus the Nijenhuis-Richardson bracket defines a Lie superalgebra structure in cohomology $H^*(\mathfrak{g}, \mathfrak{g})$, i.e., the set of bilinear functions

$$[\cdot] : H^p(\mathfrak{g}, \mathfrak{g}) \times H^q(\mathfrak{g}, \mathfrak{g}) \rightarrow H^{p+q-1}(\mathfrak{g}, \mathfrak{g})$$

with the properties (14).

Proposition 5.2. *Let $\mathfrak{g} = \oplus_{\alpha} \mathfrak{g}_{\alpha}$ be a $\mathbb{Z}$ -graded Lie algebra. Then the $\mathbb{Z}$ -grading of $C^*(\mathfrak{g}, \mathfrak{g})$ and $H^*(\mathfrak{g}, \mathfrak{g})$ is compatible with the Nijenhuis-Richardson bracket:*

$$(15) \quad \begin{aligned} [,] : C_{(\mu)}^p(\mathfrak{g}, \mathfrak{g}) \times C_{(\nu)}^q(\mathfrak{g}, \mathfrak{g}) &\longrightarrow C_{(\mu+\nu)}^{p+q-1}(\mathfrak{g}, \mathfrak{g}) \\ [,] : H_{(\mu)}^p(\mathfrak{g}, \mathfrak{g}) \times H_{(\nu)}^q(\mathfrak{g}, \mathfrak{g}) &\longrightarrow H_{(\mu+\nu)}^{p+q-1}(\mathfrak{g}, \mathfrak{g}) \end{aligned}$$

Definition 5.3. A deformation Ψ of a $\mathbb{Z}_{>0}$ -graded Lie algebra $\mathfrak{g} = \oplus_{\alpha>0} \mathfrak{g}_{\alpha}$ is called $\mathbb{Z}_{>0}$ -filtered if the following condition holds:

$$\forall X_1 \in \mathfrak{g}_{\alpha_1}, \dots, \forall X_q \in \mathfrak{g}_{\alpha_q}, \quad \Psi(X_1, \dots, X_q) \in \bigoplus_{\alpha > \alpha_1 + \alpha_2 + \dots + \alpha_q} V_{\alpha}.$$

Or in other words:

$$(16) \quad \Psi = \Psi_1 + \Psi_2 + \dots + \Psi_i + \dots, \quad \Psi_i \in C_{(i)}^2(\mathfrak{g}, \mathfrak{g}), \quad i = 1, 2, \dots$$

Decomposing $\Psi = \Psi_1 + \Psi_2 + \dots$ in the deformation equation (11) and comparing terms with the same grading we came to the following system of equations on homogeneous components Ψ_l :

$$(17) \quad \begin{aligned} d\Psi_1 = 0, \quad d\Psi_2 + \frac{1}{2}[\Psi_1, \Psi_1] = 0, \quad d\Psi_3 + [\Psi_1, \Psi_2] = 0, \dots, \\ d\Psi_i + \frac{1}{2} \sum_{m+l=i} [\Psi_m, \Psi_l] = 0, \dots \end{aligned}$$

This is the well-known system of equations for one-parametric deformation, in our case one can associate to Ψ the following one-parametric deformation Ψ_t :

$$\Psi_t = t\Psi_1 + t^2\Psi_2 + \dots + t^i\Psi_i + \dots$$

1) the first equality of (17) shows that Ψ_1 is a cocycle, we denote by $\bar{\Psi}_1$ its cohomology class in $H_{(1)}^2(\mathfrak{g}, \mathfrak{g})$.

2) the second one shows that the Nijenhuis-Richarson product $[\bar{\Psi}_1, \bar{\Psi}_1]$ defines a trivial element in $H_{(1)}^3(\mathfrak{g}, \mathfrak{g})$. If $[\bar{\Psi}_1, \bar{\Psi}_1] = 0$ then Ψ_2 is determined not uniquely but up to some closed element in $Z_{(2)}^2(\mathfrak{g}, \mathfrak{g})$.

3) hence one can find Ψ_3 iff $[\Psi_1, \Psi_2]$ is trivial in $H_{(3)}^3(\mathfrak{g}, \mathfrak{g})$ for some choice of Ψ_2 .

The subset in $H_{(3)}^3(\mathfrak{g}, \mathfrak{g})$ formed by elements $-[\Psi_1, \Psi_2]$, where Ψ_2 is a solution in $C_{(2)}^2(\mathfrak{g}, \mathfrak{g})$ of the equation $d\Psi_2 + \frac{1}{2}[\Psi_1, \Psi_1] = 0$ is called a triple Massey product $\langle \bar{\Psi}_1, \bar{\Psi}_1, \bar{\Psi}_1 \rangle$ and it is defined iff $[\bar{\Psi}_1, \bar{\Psi}_1] = 0$.

Definition 5.4. A Massey product of the k -th order $\langle \bar{\Psi}_1, \bar{\Psi}_1, \dots, \bar{\Psi}_1 \rangle$ is a subset in $H_{(k)}^3(\mathfrak{g}, \mathfrak{g})$ formed by classes $-\frac{1}{2} \sum_{m+l=k} [\Psi_m, \Psi_l]$ where $\Psi_j, j=2, \dots, k-1$ are solutions of the first $k-1$ equations of the system (17) and Ψ_1 is a representative of $\bar{\Psi}_1$. A Massey product $\langle \bar{\Psi}_1, \bar{\Psi}_1, \dots, \bar{\Psi}_1 \rangle$ is called trivial if it contains zero.

Remark. A Massey product $\langle \bar{\Psi}_1, \bar{\Psi}_1, \dots, \bar{\Psi}_1 \rangle$ of the the k -th order is defined iff all Massey products $\langle \bar{\Psi}_1, \bar{\Psi}_1, \dots, \bar{\Psi}_1 \rangle$ of orders less than k are trivial. One can easily show that it does not depend on the choice of Ψ_1 in $\bar{\Psi}_1$. A Lie product $[\bar{\Psi}_1, \bar{\Psi}_1]$ is called a Massey product of the second order.

Proposition 5.5. *Let be Ψ_1 an element of $H_{(1)}^2(\mathfrak{g}, \mathfrak{g})$. One can construct a deformation Ψ of $\mathfrak{g}$ with a first term equal to $\Psi_1 \in H_{(1)}^2(\mathfrak{g}, \mathfrak{g})$ if and only if all Massey products $\langle \bar{\Psi}_1, \bar{\Psi}_1, \dots, \bar{\Psi}_1 \rangle$ are trivial.*

Let Ψ and $\tilde{\Psi}$ be two deformations of a Lie algebra $\mathfrak{g}$. The question is: whether they define non-isomorphic Lie algebras or not? Or does there exist a non-degenerate linear transformation $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$(18) \quad \varphi([x, y] + \tilde{\Psi}(x, y)) = [\varphi(x), \varphi(y)] + \Psi(\varphi(x), \varphi(y))$$

Let us consider an non-degenerate linear operator of the form $\varphi = id + \varphi_1$, $\varphi_1 \in C^1_{(1)}(\mathfrak{g}, \mathfrak{g})$. We have for the first term

$$\varphi^{-1}([\varphi(x), \varphi(y)] + \Psi(\varphi(x), \varphi(y))) = [x, y] + (\Psi_1 + d\varphi_1)(x, y) + \dots$$

Proposition 5.6. *Let Ψ be some $\mathbb{Z}_{>0}$ -filtered deformation of $[\cdot, \cdot]$ and Φ_1 an arbitrary cocycle in $Z^2_{(1)}(\mathfrak{g}, \mathfrak{g})$ cohomologous to Ψ_1 . Then the Lie algebra with $[\cdot, \cdot] + \Psi$ is isomorphic to some filtered deformation $[\cdot, \cdot] + \tilde{\Psi}$, $\tilde{\Psi}_1 = \Phi_1$.*

6. COMPUTATIONS

In this section we compute $H^2(\mathcal{V}_n, \mathcal{V}_n)$ for $n \geq 12$.

Let $\mathfrak{g} = \bigoplus_{\alpha > 0} \mathfrak{g}_\alpha$ be a $\mathbb{Z}_{>0}$ -graded Lie algebra. One can define a decreasing filtration $\mathcal{F}$ of $(C^*(\mathfrak{g}, \mathfrak{g}), d)$:

$$\mathcal{F}^0 C^*(\mathfrak{g}, \mathfrak{g}) \supset \dots \supset \mathcal{F}^q C^*(\mathfrak{g}, \mathfrak{g}) \supset \mathcal{F}^{q+1} C^*(\mathfrak{g}, \mathfrak{g}) \supset \dots$$

where the subspace $\mathcal{F}^q C^{p+q}(\mathfrak{g}, \mathfrak{g})$ is spanned by $p+q$ -forms c in $C^{p+q}(\mathfrak{g}, \mathfrak{g})$ such that

$$c(X_1, \dots, X_{p+q}) \in \bigoplus_{\alpha=q} \mathfrak{g}_\alpha \quad \forall X_1, \dots, X_{p+q} \in \mathfrak{g}.$$

The filtration $\mathcal{F}$ is compatible with d .

Let us consider the corresponding spectral sequence $E_r^{p,q}$:

Proposition 6.1. $E_1^{p,q} = \mathfrak{g}_q \otimes H^{p+q}(\mathfrak{g})$.

We have the following natural isomorphisms:

$$(19) \quad \begin{aligned} C^{p+q}(\mathfrak{g}, \mathfrak{g}) &= \mathfrak{g} \otimes \Lambda^{p+q}(\mathfrak{g}^*) \\ E_0^{p,q} &= \mathcal{F}^q C^{p+q}(\mathfrak{g}, \mathfrak{g}) / \mathcal{F}^{q+1} C^{p+q}(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g}_q \otimes \Lambda^{p+q}(\mathfrak{g}^*). \end{aligned}$$

Now the proof follows from the formula for the $d_0^{p,q} : E_0^{p,q} \rightarrow E_0^{p+1,q}$:

$$d_0(X \otimes f) = X \otimes df,$$

where $X \in \mathfrak{g}$, $f \in \Lambda^{p+q}(\mathfrak{g}^*)$ and df is the standard differential of the cochain complex of $\mathfrak{g}$ with trivial coefficients.

Theorem 6.2. *Let $n \geq 12$ then*

1) $\dim H^0(\mathcal{V}_n, \mathcal{V}_n) = \dim H^0_{(n)}(\mathcal{V}_n, \mathcal{V}_n) = 1$;

2) $\dim H^1(\mathcal{V}_n, \mathcal{V}_n) = 4$, namely

$$\dim H^1_{(\mu)}(\mathcal{V}_n, \mathcal{V}_n) = \begin{cases} 1, & \mu = 0, n-4, n-3, n-2; \\ 0, & \text{otherwise.} \end{cases}$$

3a) *Let $n \geq 16$, then $\dim H^2(\mathcal{V}_n, \mathcal{V}_n) = 10$, more precisely:*

$$\dim H^2_{(\mu)}(\mathcal{V}_n, \mathcal{V}_n) = \begin{cases} 2, & \mu = -2; \\ 1, & \mu = -4, -3, -1, n-11, n-10, n-9, n-8, n-7; \\ 0, & \text{otherwise.} \end{cases}$$

3b) Let $12 \leq n \leq 15$, then $\dim H^2(\mathcal{V}_n, \mathcal{V}_n) = 11$, more precisely:

$$\dim H^2_{(\mu)}(\mathcal{V}_n, \mathcal{V}_n) = \begin{cases} 2, & \mu = -2; \\ 1, & \mu = -4, -3, -1, 1, n-11, n-10, n-9, n-8, n-7; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. For the proof we will use the spectral sequence considered above. Let us recall some results from [17], namely:

- 1) the basis of $H^1(\mathcal{V}_n)$ consists of two classes $[e^1]$ and $[e^2]$.
- 2) the space $H^2(\mathcal{V}_n)$ is 3-dimensional and generated by classes

$$g_5 = [e^2 \wedge e^3], \quad g_7 = [e^2 \wedge e^5 - 3e^3 \wedge e^4], \quad [\Omega_{n+1}] = \frac{1}{2} \left[\sum_{i+j=n+1} (j-i)e^i \wedge e^j \right]$$

of weights 5, 7, $n+1$ respectively.

- 3) $H^3(\mathcal{V}_n)$ is 5-dimensional and generated by elements g_{12} , g_{15} and

$$(20) \quad [e^2 \wedge \Omega_{n+1}], [e^2 \wedge \Omega_{n+2} - ne^3 \wedge \Omega_{n+1}], \\ [e^2 \wedge \Omega_{n+3} - (n+1)e^3 \wedge \Omega_{n+2} + \frac{n(n+1)}{2}e^4 \wedge \Omega_{n+1}]$$

of weights 12, 15, $n+3$, $n+4$, $n+5$ respectively. Where

$$\Omega_{n+2} = \frac{1}{2} \sum_{\substack{i+j=n+2 \\ i,j > 1}} (j-i)e^i \wedge e^j, \quad \Omega_{n+3} = \frac{1}{2} \sum_{\substack{i+j=n+3 \\ i,j > 2}} (j-i)e^i \wedge e^j$$

are projections of non-existing de^{n+2} and de^{n+3} to $\Lambda^2(\mathcal{V}_n)$. We have also

$$(21) \quad d\Omega_{n+2} = d(de^{n+2} - ne^1 \wedge e^{n+1}) = ne^1 \wedge \Omega_{n+1}, \\ d\Omega_{n+3} = d(de^{n+3} - (n+1)e^1 \wedge e^{n+2} - (n-1)e^2 \wedge e^n) = (n+1)e^1 \wedge \Omega_{n+2} + (n-1)e^2 \wedge \Omega_{n+1}.$$

Remark. The classes g_{12} , g_{15} are generators of $H^3(L_1)$ of weights 12, 15 respectively, they arise in Gontcharova's theorem (see [9] for details), as well as g_5 , g_7 in $H^2(L_1)$ and $[e^1]$, $[e^2]$ in $H^1(L_1)$. We will need the following formula for g_{12} :

$$g_{12} = [2e^2 \wedge e^3 \wedge e^7 - 5e^2 \wedge e^4 \wedge e^6 + 20e^3 \wedge e^4 \wedge e^5].$$

Proposition 6.3. *Let $n \geq 16$, then we have only a finite number of non-trivial differentials $d_r^{j,-j}$, $d_r^{j,1-j}$, $d_r^{j,2-j}$ of the spectral sequence E_r . They are:*

$$(22) \quad d_1^{j,-j} : e_j \mapsto (1-j)e_{j+1} \otimes [e^1], \quad j \neq 1; \\ d_2^{1,1} : e_1 \mapsto e_3 \otimes [e^2], \\ d_2^{j,2-j} : e_j \otimes [\Omega_{n+1}] \mapsto \left(j-2 - \frac{(n-1)j(j-1)}{(n+1)n} \right) e_{j+2} \otimes [e^2 \wedge \Omega_{n+1}]; \\ d_3^{j,1-j} : e_j \otimes [e^2] \mapsto -\frac{1}{6}(j-3)(j^2-3j+8)e_{j+3} \otimes [e^2 \wedge e^3], \quad j \neq 3; \\ d_4^{2,-1} : e_2 \otimes [e^1] \mapsto -\frac{4}{3}e_6 \otimes [e^2 \wedge e^3]; \\ d_5^{j,2-j} : e_j \otimes [e^2 \wedge e^5 - 3e^3 \wedge e^4] \mapsto -\frac{1}{5544}(j-8)(j^2-4j+27)(j^2-13j+48)e_{j+5} \otimes g_{12}, \quad j \neq 8; \\ d_8^{8,-6} : \text{Span}(e_8 \otimes [e^2 \wedge e^5 - 3e^3 \wedge e^4]) \rightarrow \text{Span}(e_{16} \otimes g_{15}).$$

If $12 \leq n \leq 15$ then only the differential $d_8^{8,-6}$ becomes trivial.

Corollary 6.4. *The following classes in $E_1^{p,q}, p+q=0, 1, 2$ survive to E_∞ for $n \geq 16$:*

$$(23) \quad \begin{aligned} & e_n; e_1 \otimes [e^1], e_n \otimes [e^2], e_{n-1} \otimes [e^2], e_{n-2} \otimes [e^2]; \\ & e_{n-1} \otimes [\Omega_{n+1}], \quad e_n \otimes [\Omega_{n+1}], \\ & e_i \otimes [e^2 \wedge e^3], \quad i = 1, 2, 3; \\ & e_j \otimes [e^2 \wedge e^5 - 3e^3 \wedge e^4], \quad j = n-4, n-3, n-2, n-1, n. \end{aligned}$$

If $12 \leq n \leq 15$ then one have to add $e_8 \otimes [e^2 \wedge e^5 - 3e^3 \wedge e^4]$ to the list above.

Remark. 1) the element e_n spans the center $Z(\mathcal{V}_n) = H^0(\mathcal{V}_n, \mathcal{V}_n)$;

2) the class $e_1 \otimes [e^1]$ corresponds to the inner derivation $\sum_{i=1} ie_i \otimes e^i = ad(e_0)$ in the solvable Lie algebra L_0/L_{n+1} restricted to the nilpotent ideal $\mathcal{V}_n$.

Hence the proof of our theorem follows from Corollary 6.4. It's time to say that the Proposition 6.3 it is finite-dimensional version of the following theorem

Theorem 6.5 (A. Fialowski, [7], [8]). $\dim H^2(L_1, L_1) = 3$, more precisely:

$$\dim H_{(\mu)}^2(L_1, L_1) = \begin{cases} 1, & \mu = -2, -3, -4; \\ 0, & \text{otherwise.} \end{cases}$$

The differentials $d_1^{j,-j}, d_2^{1,1}, d_3^{j,1-j}, d_4^{2,-1}, d_5^{j,2-j}, d_8^{8,-6}$ of Proposition 6.3 came from the corresponding spectral sequence for $H^*(L_1, L_1)$. Their non-triviality follows from the more general result of B. Feigin, D. Fuchs [6]. But it is a very complicate task to follow all details of the proof in [6]. So it appears to be usefull to calculate $d_1^{j,-j}, d_3^{j,1-j}, d_5^{j,2-j}$ explicitly. From the other hand explicit formulae will give us a possibility to write down structure relations of all deformations of $\mathcal{V}_n$.

Remark. In the infinite-dimensional case only classes $e_i \otimes [e^2 \wedge e^3], i = 1, 2, 3$ survive to E_∞ and they correspond to generators in $H^2(L_1, L_1)$ of weights $\mu = i - (2+3) = -4, -3, -2$ that were found by A. Fialowski.

The proof of Proposition 6.3 consists of direct calculations.

First of all by definiton of $d: C^0(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \rightarrow C^1(\mathfrak{g}, \mathfrak{g}) = \text{Hom}(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes \mathfrak{g}^*$ we have

$$d(e_j) = (j-1)e_{j+1} \otimes e^1 + (j-2)e_{j+2} \otimes e^2 + (j-3)e_{j+3} \otimes e^3 + \dots$$

Hence

$$d_1^{j,-j}(e_j) = e_{j+1} \otimes (j-1)[e^1], \quad j \neq 1.$$

Now we go to the differential d_2 .

$$d(e_j \otimes \Omega_{n+1}) = e_{j+1} \otimes (j-1)e^1 \wedge \Omega_{n+1} + \dots$$

where dots stand instead of terms of higher filtration. As we know $e^1 \wedge \Omega_n = \frac{1}{n} d\Omega_{n+2}$ and we now can take new representative $e_j \otimes \Omega_n + \frac{j-1}{n} e_{j+1} \otimes \Omega_n$ such that:

$$d(e_j \otimes \Omega_{n+1} + \frac{j-1}{n} e_{j+1} \otimes \Omega_{n+2}) = e_{j+2} \otimes \left(\frac{j(j-1)}{n} e^1 \wedge \Omega_{n+2} + (j-2)e^2 \wedge \Omega_{n+1} \right) + \dots$$

But $d\Omega_{n+3} = (n+1)e^1 \wedge \Omega_{n+2} + (n-1)e^2 \wedge \Omega_{n+1}$ and therefore $\frac{j(j-1)}{n} e^1 \wedge \Omega_{n+2} + (j-2)e^2 \wedge \Omega_{n+1}$ is cohomologous to $\left(j-2 - \frac{(n-1)j(j-1)}{(n+1)n} \right) e^2 \wedge \Omega_{n+1}$. Hence we conclude

$$d_2(e_j \otimes [\Omega_{n+1}]) = e_{j+2} \otimes \left(j-2 - \frac{(n-1)j(j-1)}{(n+1)n} \right) [e^2 \wedge \Omega_{n+1}].$$

Now the most complicated case: computation of $d_2(e_j \otimes [e^2 \wedge e^5 - 3e^3 \wedge e^4])$. We have to find $\xi_1, \xi_2, \xi_3, \xi_4$ such that

$$(24) \quad d(e_j \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4)) + \sum_{p=1}^4 e_{j+p} \otimes \xi_p = e_{j+5} \otimes \xi + \dots$$

for some $\xi \in C^3(\mathcal{V}_n, \mathcal{V}_n)$. We recall the notation $g_7 = e^2 \wedge e^5 - 3e^3 \wedge e^4$. We have the following system of equations on $\xi_1, \xi_2, \xi_3, \xi_4$:

$$(25) \quad \begin{aligned} d\xi_1 &= (j-1)e^1 \wedge g_7; \\ d\xi_2 &= je^1 \wedge \xi_1 + (j-2)e^2 \wedge g_7; \\ d\xi_3 &= (j+1)e^1 \wedge \xi_2 + (j-2)e^2 \wedge \xi_1 + (j-3)e^3 \wedge g_7; \\ d\xi_4 &= (j+2)e^1 \wedge \xi_3 + je^2 \wedge \xi_2 + (j-2)e^2 \wedge \xi_1 + (j-4)e^4 \wedge g_7. \end{aligned}$$

Taking $\xi_1, \xi_2, \xi_3, \xi_4$ as homogeneous 2-forms of weights 8, 9, 10, 11 one can remark that the right parts of these equations are exact forms because $H_{(p)}^3(\mathcal{V}_n) = 0, p \leq 11$. $\xi_1, \xi_2, \xi_3, \xi_4$ are defined uniquely by the condition $\xi_p \in \Lambda^2(e^2, \dots, e^n), p = 1, 2, 3, 4$.

The answer is:

$$(26) \quad \begin{aligned} \xi_1 &= \frac{j-1}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5); \\ \xi_2 &= P_1(j)e^2 \wedge e^7 + P_2(j)e^3 \wedge e^6 + P_3(j)e^4 \wedge e^5; \\ \xi_3 &= Q_1(j)e^2 \wedge e^8 + Q_2(j)e^3 \wedge e^7 + Q_3(j)e^4 \wedge e^6; \\ \xi_4 &= Z_1(j)e^2 \wedge e^9 + Z_2(j)e^3 \wedge e^8 + Z_3(j)e^4 \wedge e^7 + Z_4(j)e^5 \wedge e^6, \end{aligned}$$

where polinomials $P_i(j), Q_j(j), Z_l(j)$ are defined by

$$(27) \quad \begin{aligned} P_1(j) &= \frac{(5\binom{j}{2} + 3\binom{j}{1} - 6)}{21}; \quad P_2(j) = -\frac{(4\binom{j}{2} + 15\binom{j}{1} - 30)}{21}; \quad P_3(j) = -\frac{(13\binom{j}{2} - 30\binom{j}{1} + 60)}{21}; \\ Q_1(j) &= \frac{(3\binom{j+1}{3} + 4\binom{j+1}{2} - 4\binom{j+1}{1})}{28}; \quad Q_2(j) = \frac{((\binom{j+1}{3}) - 8\binom{j+1}{2} + 8\binom{j+1}{1})}{14}; \\ Q_3(j) &= \frac{(-13\binom{j+1}{3} + 20\binom{j+1}{2} - 20\binom{j+1}{1})}{28}; \\ Z_1(j) &= \frac{1}{22}\binom{j+2}{4} + \frac{23}{231}\binom{j+2}{3} - \frac{7}{198}\binom{j+2}{2} - \frac{59}{693}\binom{j+2}{1} + \frac{37}{231}; \\ Z_2(j) &= \frac{17}{154}\binom{j+2}{4} - \frac{62}{231}\binom{j+2}{3} - \frac{53}{1386}\binom{j+2}{2} - \frac{59}{99}\binom{j+2}{1} - \frac{37}{33}; \\ Z_3(j) &= -\frac{29}{154}\binom{j+2}{4} - \frac{4}{77}\binom{j+2}{3} + \frac{317}{462}\binom{j+2}{2} - \frac{59}{33}\binom{j+2}{1} + \frac{37}{11}; \\ Z_4(j) &= -\frac{47}{154}\binom{j+2}{4} + \frac{185}{231}\binom{j+2}{3} - \frac{2245}{1386}\binom{j+2}{2} + \frac{295}{99}\binom{j+2}{1} - \frac{185}{33}; \end{aligned}$$

Now one can calculate ξ from equation (24):

$$\xi = (j+3)e^1 \wedge \xi_4 + (j+1)e^2 \wedge \xi_3 + (j-1)e^3 \wedge \xi_2 + (j-3)e^4 \wedge \xi_1 + (j-5)e^5 \wedge g_7.$$

The space $H_{(12)}^3(\mathcal{V}_n)$ is one-dimensional and we have for the cohomology class $[\xi]$:

$$[\xi] = -\frac{1}{5544}(j-8)(j^2-4j+27)(j^2-13j+48) [2e^2 \wedge e^3 \wedge e^7 - 5e^2 \wedge e^4 \wedge e^6 + 20e^3 \wedge e^4 \wedge e^5].$$

In the same way one can remark that

$$(28) \quad \begin{aligned} & d(e_j \otimes e^2 + e_{j+1} \otimes (j-1)e^3 + e_{j+2} \otimes \frac{j(j-1)}{2}e^4) = \\ & = e_{j+3} \otimes \left(\frac{(j+1)j(j-1)}{2}e^1 \wedge e^4 + ((j-1)^2 - (j-3))e^2 \wedge e^3 \right) + \dots \end{aligned}$$

As $3e^1 \wedge e^4$ is cohomologous to $e^2 \wedge e^3$ it follows that

$$d_3(e_j \otimes [e^2]) = e_{j+3} \otimes \left((j-1)^2 - (j-3) - \frac{(j+1)j(j-1)}{6} \right) [e^2 \wedge e^3]$$

□

Now we choose the following basic cocycles $\psi_{k,i}$ in $\oplus_{\mu>0} H_{(\mu)}^2(\mathcal{V}_n, \mathcal{V}_n)$, $n \geq 16$:

	basic cocycles
$H_{(n-7)}^2(\mathcal{V}_n, \mathcal{V}_n)$	$\psi_{n,7} = e_n \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4)$
$H_{(n-8)}^2(\mathcal{V}_n, \mathcal{V}_n)$	$\psi_{n,8} = e_{n-1} \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4) + e_n \otimes \frac{n-2}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5);$
$H_{(n-9)}^2(\mathcal{V}_n, \mathcal{V}_n)$	$\psi_{n,9} = e_{n-2} \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4) + e_{n-1} \otimes \frac{n-3}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5) + e_n \otimes (P_1(n-2)e^2 \wedge e^7 + P_2(n-2)e^3 \wedge e^6 + P_3(n-2)e^4 \wedge e^5)$
$H_{(n-10)}^2(\mathcal{V}_n, \mathcal{V}_n)$	$\psi_{n,10} = e_{n-3} \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4) + e_{n-2} \otimes \frac{n-4}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5) + e_{n-1} \otimes (P_1(n-3)e^2 \wedge e^7 + P_2(n-3)e^3 \wedge e^6 + P_3(n-3)e^4 \wedge e^5) + e_n \otimes (Q_1(n-3)e^2 \wedge e^8 + Q_2(n-3)e^3 \wedge e^7 + Q_3(n-3)e^4 \wedge e^6)$
$H_{(n-11)}^2(\mathcal{V}_n, \mathcal{V}_n)$	$\psi_{n,11} = e_{n-4} \otimes (e^2 \wedge e^5 - 3e^3 \wedge e^4) + e_{n-3} \otimes \frac{n-5}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5) + e_{n-2} \otimes (P_1(n-4)e^2 \wedge e^7 + P_2(n-4)e^3 \wedge e^6 + P_3(n-4)e^4 \wedge e^5) + e_{n-1} \otimes (Q_1(n-4)e^2 \wedge e^8 + Q_2(n-4)e^3 \wedge e^7 + Q_3(n-4)e^4 \wedge e^6) + e_n \otimes (Z_1(n-4)e^2 \wedge e^9 + Z_2(n-4)e^3 \wedge e^8 + Z_3(n-4)e^4 \wedge e^7 + Z_4(n-4)e^5 \wedge e^6)$

7. MODULI SPACE OF $\mathbb{Z}_{>0}$ -FILTERED DEFORMATIONS.

In this section we classify up to an isomorphism the Lie algebras over $\mathbb{K}$ defined by the basis $e_1, \dots, e_n$, $n \geq 16$ and commuting relations of the following form:

$$(29) \quad [e_i, e_j] = (j-i)e_{i+j} + \sum_{l=1}^{n-i-j} c_{ij}^l e_{i+j+l}.$$

A Lie algebra $\mathfrak{g}$ with the commuting relations (29) is a $\mathbb{Z}_{>0}$ -filtered deformation of the $\mathbb{Z}_{>0}$ -graded Lie algebra $\mathcal{V}_n$, i.e. $\mathfrak{g} = (\mathcal{V}_n, [,] + \Psi)$, where

$$(30) \quad \begin{aligned} & \Psi = \Psi_1 + \Psi_2 + \dots + \Psi_{n-3}, \quad \Psi_l \in C_{(l)}^2(\mathcal{V}_n, \mathcal{V}_n), \\ & \Psi_l(e_i, e_j) = \begin{cases} c_{ij}^l e_{i+j+l}, & i+j \leq n-l, \\ 0, & \text{otherwise;} \end{cases} \quad l = 1, 2, \dots, n-3. \end{aligned}$$

satisfying to the system (17) of deformation equations:

$$d\Psi_1 = 0, \quad d\Psi_2 + \frac{1}{2}[\Psi_1, \Psi_1] = 0, \quad \dots, \quad d\Psi_{n-6} + \frac{1}{2} \sum_{i+j=n-6} [\Psi_i, \Psi_j] = 0.$$

Lemma 7.1. *Let $\Psi, \tilde{\Psi}$ be two $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n, n \geq 5$ and*

$$\varphi : \mathfrak{g} = (\mathcal{V}_n, [,] + \Psi) \rightarrow \tilde{\mathfrak{g}} = (\mathcal{V}_n, [,] + \tilde{\Psi})$$

is a Lie algebra isomorphism.

Then

$$(31) \quad \begin{aligned} 1) & \quad \varphi = \varphi_0 + \varphi_1 + \varphi_2 + \dots + \varphi_{n-1}, \quad \varphi_j \in C_{(j)}^1(\mathcal{V}_n, \mathcal{V}_n), \quad j = 0, 1, 2, \dots, n-1, \\ 2) & \quad \varphi_0(e_i) = \alpha^i e_i, \quad \alpha \in \mathbb{K}^*, \quad i = 1, 2, \dots, n. \end{aligned}$$

Proof. We will study the dual situation. The mapping

$$\varphi^* : \mathfrak{g}^* = (\mathcal{V}_n^*, d_\Psi = d + \Psi^*) \rightarrow \tilde{\mathfrak{g}}^* = (\mathcal{V}_n^*, d_{\tilde{\Psi}} = d + \tilde{\Psi}^*)$$

is an isomorphism of d -algebras.

For the dual basis $e^1, e^2, \dots, e^n$ of $(\mathcal{V}_n^*, d_\Psi)$ we have the following structure relations:

$$d_\Psi e^k = \frac{1}{2} \sum_{i+j=k} (j-i) e^i \wedge e^j + \frac{1}{2} \sum_{m+p < k} c_{mp}^k e^m \wedge e^p, \quad k = 3, \dots, n.$$

Let us write down some of them.

$$(32) \quad \begin{aligned} d_\Psi e^1 &= d_\Psi e^2 = 0, \quad d_\Psi e^3 = e^1 \wedge e^2, \quad d_\Psi e^4 = 2e^1 \wedge e^3 + c_{12}^1 e^1 \wedge e^2, \\ d_\Psi e^5 &= 3e^1 \wedge e^4 + e^2 \wedge e^3 + c_{13}^1 e^1 \wedge e^3 + c_{12}^2 e^1 \wedge e^2, \dots \end{aligned}$$

The dual mapping $\varphi^* : \tilde{\mathfrak{g}}^* \rightarrow \mathfrak{g}^*$ is the isomorphism of d -algebras, i.e.

$$d_\Psi \varphi^* = \varphi^* d_{\tilde{\Psi}}$$

$$1) \quad d_\Psi \varphi^* e^1 = d_\Psi \varphi^* e^2 = 0, \text{ thus}$$

$$\varphi^* e^1 = \alpha_{11} e^1 + \alpha_{21} e^2; \quad \varphi^* e^2 = \alpha_{12} e^1 + \alpha_{22} e^2.$$

$$2) \quad d_\Psi \varphi^* e^3 = \varphi^* e^1 \wedge \varphi^* e^2 = (\alpha_{11} \alpha_{22} - \alpha_{21} \alpha_{12}) e^1 \wedge e^2, \text{ and we have}$$

$$\varphi^* e^3 = (\alpha_{11} \alpha_{22} - \alpha_{21} \alpha_{12}) e^3 + \alpha_{13} e^1 + \alpha_{23} e^2.$$

3) The form $\varphi^* d_{\tilde{\Psi}} e^4$ is cohomologous to $2\alpha_{21}(\alpha_{11} \alpha_{22} - \alpha_{21} \alpha_{12}) e^2 \wedge e^3$ and it is exact iff $\alpha_{21} = 0$, thus

$$\varphi^* e^4 = \alpha_{11}^2 \alpha_{22} e^4 + (\alpha_{11} \alpha_{23} + \tilde{c}_{12}^1 \alpha_{11} \alpha_{22} - c_{12}^1 \alpha_{11}^2 \alpha_{22}) e^3 + \alpha_{14} e^1 + \alpha_{24} e^2.$$

4) $\varphi^* d_{\tilde{\Psi}} e^5 \sim 3\alpha_{22} \alpha_{11}^3 e^1 \wedge e^4 + \alpha_{22}^2 \alpha_{11} e^2 \wedge e^3$ and the last one is exact iff $\alpha_{12}^2 = \alpha_{22}$, and for the moment we have

$$(33) \quad \begin{aligned} \varphi^* e^1 &= \alpha_{11} e^1, \quad \varphi^* e^2 = \alpha_{11}^2 e^2 + \alpha_{12} e^1, \\ \varphi^* e^3 &= \alpha_{11}^3 e^3 + \alpha_{13} e^1 + \alpha_{23} e^2, \\ \varphi^* e^4 &= \alpha_{11}^4 e^4 + \dots, \quad \varphi^* e^5 = \alpha_{11}^5 e^5 + \dots \end{aligned}$$

Going on and using an obvious inductive assumption we have for the operator φ^* :

$$(34) \quad \varphi^* e^i = \alpha^i e^i + \sum_{l < i}^n \alpha_{li} e^l, \quad i = 1, \dots, n,$$

for some $\alpha \neq 0, \alpha_{li} \in \mathbb{K}$. □

From the other hand it is evident that changing the canonical basis of an arbitrary Lie algebra $\mathfrak{g}$ of the type (29) by an operator φ with property (34) we will get again the commuting relations of the same type.

Corollary 7.2. *The matrix Lie group G_n of lower-triangular matrixes φ of the following type*

$$\varphi = \begin{pmatrix} \alpha & 0 & \dots & 0 \\ a_{21} & \alpha^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{nn-1} & \dots & \alpha^n \end{pmatrix}, \quad a_{ij}, \alpha \in \mathbb{K}, \alpha \neq 0,$$

acts on the set V_n of $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$ as the group of changes of canonical basis:

$$(35) \quad (\varphi \star \Psi)(x, y) = \varphi^{-1}([\varphi x, \varphi y] + \Psi(\varphi x, \varphi y)) - [x, y], \quad \forall x, y \in \mathcal{V}_n, \varphi \in G_n.$$

Two $\mathbb{Z}_{>0}$ -filtered deformations $(\mathcal{V}_n, [\cdot, \cdot] + \Psi)$ and $(\mathcal{V}_n, [\cdot, \cdot] + \tilde{\Psi})$ are isomorphic as Lie algebras if and only if they are in the same orbit O_Ψ of the G_n -action.

Hence we have proved the following

Theorem 7.3. *Let $n \geq 5$, then there is a one-to-one correspondance between the orbit space $O(G_n, V_n)$ of the action G_n on the set V_n of $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$ and the moduli space $\mathcal{M}_n$ (the set of isomorphism classes of the $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$).*

Proposition 7.4. *The matrix Lie group G_n is the semi-direct product $\mathbb{K}^* \ltimes UT_n$ where UT_n denotes the group of unitriangular matrixes:*

$$G_n = \mathbb{K}^* \ltimes UT_n = \mathbb{K}^* \ltimes \left\{ \begin{pmatrix} 1 & 0 & \dots & 0 \\ * & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \dots & 1 \end{pmatrix} \right\}, \quad \mathbb{K}^* \cong \left\{ \begin{pmatrix} \alpha & 0 & \dots & 0 \\ 0 & \alpha^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \alpha^n \end{pmatrix} \right\}.$$

Remark (see also section 5). If $\Psi = \Psi_1 + \Psi_2 + \dots + \Psi_{n-3}$ is a solution of the system (17) of deformation equations then $\Psi = t\Psi_1 + t^2\Psi_2 + \dots + t^{n-3}\Psi_{n-3}$, $\forall t \in \mathbb{K}$ also satisfies to the system (17). It follows that the space V_n can be retracted over itself to $\mathcal{V}_n$. From another hand one can define $\mathbb{K}^*$ -action on V_n :

$$\rho_n(\alpha)(\Psi_1, \Psi_2, \dots, \Psi_{n-3}) = (\alpha\Psi_1, \alpha^2\Psi_2, \dots, \alpha^{n-3}\Psi_{n-3}), \quad \alpha \in \mathbb{K}^*.$$

Evidently this action coincides with the action on V_n of the subgroup $\mathbb{K}^*$ of diagonal matrixes in G_n .

Proposition 7.5. *There are the following bijections:*

$$O_{\rho_n}(\mathbb{K}^*, O(UT_n, V_n)) \rightarrow O(G_n/UT_n, O(UT_n, V_n)) \rightarrow O(G_n, V_n).$$

Proposition 7.6. *Let Ψ be a $\mathbb{Z}_{>0}$ -filtered deformation of $\mathcal{V}_n$, $n \geq 12$. Then there exists an element $\tilde{\Psi}$ in the UT_n -orbit O_Ψ of Ψ such that*

$$\tilde{\Psi}_1 = \dots = \tilde{\Psi}_{n-12} = 0.$$

For an arbitrary $\mathbb{Z}_{>0}$ -filtered deformation Ψ the first equation is $d\Psi_1 = 0$. We recall now

$$H_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) = 0, \quad i \leq n-12$$

and hence $\tilde{\Psi}_1 = d\varphi_1$ for some φ_1 of $C_{(1)}^1(\mathcal{V}_n, \mathcal{V}_n)$. Acting by $g = id + \varphi_1 \in UT_n$ we get $\tilde{\Psi} = g \star \Psi$ such that $\tilde{\Psi}_1 = 0$. Now the second equation of the system for this new element $\tilde{\Psi}$ will be $d\tilde{\Psi}_2 = [\tilde{\Psi}_1, \tilde{\Psi}_1] = 0$. We act on $\tilde{\Psi}$ by $g = id + \varphi_2$, where $\tilde{\Psi}_2 = d\varphi_2$ and continue the procedure step by step.

Now we suppose to be constructed an element $\tilde{\Psi}$ such that $\tilde{\Psi}_1 = \dots = \tilde{\Psi}_{n-12} = 0$ satisfying

$$d\tilde{\Psi}_{n-11} = 0.$$

We recall that in the section 6 we found the basic cocycles $\psi_{n,i}$, such that

$$\text{Span}([\psi_{n,i}]) = H_{(n-i)}^2(\mathcal{V}_n, \mathcal{V}_n), \quad i = 7, \dots, 11.$$

Hence $\tilde{\Psi}_{n-11} = x_1\psi_{n,11} + d\varphi_{n-11}$, $x_1 \in \mathbb{K}$, and $\varphi_{n-11} \in C_{(n-11)}^1(\mathcal{V}_n, \mathcal{V}_n)$. Then acting on $\tilde{\Psi}$ by $id + \varphi_{n-11}$ we get again a new element in the orbit O_Ψ (we keep the same notation $\tilde{\Psi}$ for it) with the property $\tilde{\Psi}_{n-11} = x_1\psi_{n,11}$.

Proposition 7.7. *Let $n \geq 14$ then all Nijenhuis-Richardson products of basic cocycles $\psi_{n,i}$ are trivial elements in $C^3(\mathcal{V}_n, \mathcal{V}_n)$:*

$$[\psi_{n,i}, \psi_{n,j}](x, y, z) = 0 \quad \forall x, y, z \in \mathcal{V}_n.$$

The proof follows from the two properties of $\psi_{n,i}$:

- 1) $\text{Im}\psi_{n,i} = \text{Span}(e_n, \dots, e_{n-4})$;
- 2) $\psi_{n,i}(x, y) = 0$ if $x \wedge y \notin \Lambda^2(e_2, \dots, e_9)$.

Hence $\psi_{n,i}(\psi_{n,j}(x, y), z) = 0$, $\forall x, y, z$ if $n-4 > 9$.

Proposition 7.8. *Let $n \geq 14$. There is a one-to-one correspondance between the orbit space $O(UT_n, V_n)$ and the 5-dimensional vector space $\oplus_{i>0} H_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$.*

This proposition follows from the previous one. Namely in an arbitrary UT_n -orbit O_Ψ one can choose the unique representative $\tilde{\Psi}$ such that

$$\tilde{\Psi} = x_1\psi_{n,11} + x_2\psi_{n,10} + x_3\psi_{n,9} + x_4\psi_{n,8} + x_5\psi_{n,7},$$

where $x_i \in \mathbb{K}, i = 1, \dots, 5$. We will call $\tilde{\Psi}$ the canonical element of the orbit O_Ψ and the set $\{x_1, \dots, x_5\}$ is called the homogeneous coordinates of the orbit $O_\Psi \in O(UT_n, V_n)$.

Theorem 7.9. *Let $n \geq 16$. There is a one-to-one correspondance between the moduli space $\mathcal{M}_n = O(G_n, V_n)$ of $\mathbb{Z}_{>0}$ -filtered deformations of $\mathcal{V}_n$ and the orbit space $O_{\tilde{\rho}_n}(\mathbb{K}^*, \mathbb{K}^5)$ where the action $\tilde{\rho}_n$ of $\mathbb{K}^*$ on $\mathbb{K}^5$ is defined in coordinates x^i :*

$$(36) \quad \tilde{\rho}_n(\alpha)(x_1, x_2, \dots, x_5) = (\alpha^{n-11}x_1, \alpha^{n-10}x_2, \dots, \alpha^{n-7}x_5), \quad \alpha \in \mathbb{K}^*.$$

We have to verify only the formula for $\mathbb{K}^*$ -action on $O(UT_n, V_n)$. But

$$(37) \quad \begin{aligned} \tilde{\rho}_n(\alpha)(x_1, x_2, \dots, x_5) &= \rho_n(\alpha)(x_1\psi_{n,11} + x_2\psi_{n,10} + \dots + x_5\psi_{n,7}) = \\ &= \alpha^{n-11}x_1\psi_{n,11} + \alpha^{n-10}x_2\psi_{n,10} + \dots + \alpha^{n-7}x_5\psi_{n,7}. \end{aligned}$$

8. AFFINE VARIETY OF $\mathbb{Z}_{>0}$ -FILTERED DEFORMATIONS

One can regard the set V_n of $\mathbb{Z}_{>0}$ -filtered deformation of $\mathcal{V}_n$ as a affine variety in the affine space $\oplus_{i>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$. Namely one can define a mapping (polynomial in the coordinates $\{c_{ij}^k\}$):

$$\mathcal{F} : \oplus_{i>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{i>0} C_{(i)}^3(\mathcal{V}_n, \mathcal{V}_n),$$

where

$$\mathcal{F}(\Psi_1, \Psi_2, \dots, \Psi_{n-3}) = (d\Psi_1, d\Psi_2 + \frac{1}{2}[\Psi_1, \Psi_1], \dots, d\Psi_{n-6} + \frac{1}{2} \sum_{i+j=n-6} [\Psi_i, \Psi_j]).$$

Then

$$V_n = \{\Psi \in \oplus_{i>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \mid \mathcal{F}(\Psi) = 0\}.$$

Theorem 8.1. (1) *The variety V_n has no singular points and*

$$\dim V_n = \begin{cases} \frac{n(n-3)}{2} + 3, & n \geq 16; \\ \frac{n(n-3)}{2} + 4, & 12 \leq n \leq 15. \end{cases}$$

(2) *The group UT_n acts on V_n with constant rank and we have for the dimension of an arbitrary orbit O_Ψ*

$$\dim O_\Psi = \frac{n(n-3)}{2} - 2.$$

One can identify the Lie algebra of UT_n with $\oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n)$. Let Ψ be some fixed element of V_n , then decomposing $\varphi = \exp \alpha = 1 + \alpha + \alpha^2 + \dots$ in the formula (35) and taking linear terms with respect to α one can get the following formula for the differential D of the UT_n -action ($x, y \in \mathcal{V}_n$):

$$D_\Psi(\alpha)(x, y) = [\alpha(x), y] + [x, \alpha(y)] - \alpha([x, y]) + \Psi(\alpha(x), y) + \Psi(x, \alpha(y)) - \alpha(\Psi(x, y)).$$

Hence

$$D_\Psi(\alpha) = -d(\alpha) + [\Psi, \alpha] = -d_\Psi(\alpha), \quad \alpha \in \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n)$$

where $d_\Psi : \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$ is a deformed differential of the cochain complex of $\mathcal{V}_n$. But in the same time d_Ψ defines the differential of the cochain complex of the Lie algebra $\mathfrak{g} = (\mathcal{V}_n, [,] + \Psi)$.

Analogously one can show that the differential $D\mathcal{F}_\Psi$ coincides with the differential $d_\Psi : \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^3(\mathcal{V}_n, \mathcal{V}_n)$ of the cochain complex with coefficients in the adjoint representation of $\mathfrak{g} = (\mathcal{V}_n, [,] + \Psi)$.

Remark. Let us denote by $d_i : C_{(i)}^*(\mathcal{V}_n, \mathcal{V}_n) \rightarrow C_{(i)}^{*+1}(\mathcal{V}_n, \mathcal{V}_n)$ the restriction of the differential d of the cochain complex $(C^*(\mathcal{V}_n, \mathcal{V}_n), d)$ to homogeneous components. Then the matrix of d_Ψ for an arbitrary Ψ is a block-triangular for some natural number l :

$$d_\Psi = \begin{pmatrix} d_1 & * & \dots & * \\ 0 & d_2 & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_{n-l} \end{pmatrix}$$

And we have the following estimates

$$\text{rank } d_\Psi \geq \text{rank } d, \quad \dim \text{Ker } d_\Psi \leq \dim \text{Ker } d.$$

In order to compute these ranks one may assume that $\Psi = \tilde{\Psi}$, where $\tilde{\Psi} \in \mathbb{K}^5$ is the canonical representative of the orbit O_Ψ .

Proposition 8.2.

$$\begin{aligned} (38) \quad & \dim \text{Ker} \left(d_{\tilde{\Psi}} : \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \right) = \\ & = \dim \text{Ker} \left(d : \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \right) = n + 2. \end{aligned}$$

One can easily verify that operators

$$(39) \quad \begin{aligned} & ad_{\tilde{\Psi}}(e_1), ad_{\tilde{\Psi}}(e_2), \dots, ad_{\tilde{\Psi}}(e_{n-1}), e_n \otimes e^2, e_{n-1} \otimes e^2 + (n-2)e_n \otimes e^3, \\ & e_{n-2} \otimes e^2 + (n-3)e_n \otimes e^3 + \frac{(n-2)(n-3)}{2} e_n \otimes e^3 \end{aligned}$$

give the basis of $\text{Ker } d_{\tilde{\Psi}} : \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n)$.

Hence

$$\dim O_{\Psi} = \dim \text{Im} \left(d_{\Psi} : \oplus_{>0} C_{(i)}^1(\mathcal{V}_n, \mathcal{V}_n) \rightarrow \oplus_{>0} C_{(i)}^2(\mathcal{V}_n, \mathcal{V}_n) \right) = \frac{n(n-1)}{2} - n - 2.$$

From the other hand the cocycles $\{\psi_{n,i} | i=7, \dots, 11\}$ span the tangent space to $\mathbb{K}^5$ in an arbitrary $\tilde{\Psi}$, moreover $\{\psi_{n,i}\}$ are linearly independent modulo $\text{Im } d_{\Psi}$ because $\{[\psi_{n,i}] | i=7, \dots, 11\}$ is the basis in $\oplus_{>0} H_{(i)}^2(\mathfrak{g}, \mathfrak{g})$.

The variety V_n can be regarded as a subvariety of the affine variety $\mathcal{N}_n$ of n -dimensional nilpotent Lie algebras. There exists a GL_n -action on $\mathcal{N}_n$ by basis changes. Let us consider the set $V'_n = \{y \in \mathcal{N}_n | \exists \Psi \in V_n, \exists g \in GL_n, y = g \star \Psi\}$.

Remark. In fact the Zariski closure of V'_n in $\mathcal{N}_n$ coincides with one of irreducible components of $\mathcal{N}_n$ that were discussed by Yu.Khakimdjanov in [13].

The GL_n -action on $V'_n = \{y \in \mathcal{N}_n | \exists x \in V_n, \exists g \in GL_n, y = g \star x\}$ has singularities. Namely by the lemma 7.1 the stabilizer $(GL_n)_{\tilde{\Psi}}$ of a point $\tilde{\Psi} \in \mathbb{K}^5 \subset V_n$ coincides with the stabilizer $(G_n)_{\tilde{\Psi}}$ and we have

$$\dim(GL_n)_{\tilde{\Psi}} = \begin{cases} n+2, & \text{if } \tilde{\Psi} \neq 0; \\ n+3, & \text{if } \tilde{\Psi} = 0. \end{cases}$$

Remark. If $\tilde{\Psi} = 0$ then one have to add the operator $\sum_{i=1}^n i e_i \otimes e^i$ to (39) in order to get a basis of the Lie algebra of the stabilizer $(GL_n)_{\tilde{\Psi}}$.

Moreover, even for non-zero points $\tilde{\Psi}$ there are some particularities. Let $\tilde{\Psi}$ be a generic point in $\mathbb{K}^5$, i.e. $x_i \neq 0, \forall i$, then

$$(GL_n)_{\tilde{\Psi}} = (G_n)_{\tilde{\Psi}} = (UT_n)_{\tilde{\Psi}}.$$

But if $\mathbb{K} = \mathbb{C}$ then for instance for $\tilde{\psi} = (1, 0, 0, 0, 0)$ we have

$$(GL_n)_{\tilde{\Psi}} = (G_n)_{\tilde{\Psi}} = \mathbb{Z}_{n-11} \ltimes (UT_n)_{\tilde{\Psi}},$$

where $\mathbb{Z}_{n-11}$ stands for the group of the roots of the unit:

$$\mathbb{Z}_{n-11} = \{\alpha \in \mathbb{C} \mid \alpha^{n-11} = 1\}.$$

It is interesting here to remark some parallels between our discussions and the theory of non-singular deformations, that was introduced in [8].

Definition 8.3 ([8]). Let $\mathfrak{g}$ be a Lie algebra with the comutator $[\cdot, \cdot]$. Consider a formal one-parameter deformation

$$[x, y]_t = [x, y] + \sum_{k \geq 1} \alpha_k(x, y) t^k$$

of $\mathfrak{g}$. A deformation is called non-singular if there exists a formal one-parameter family of linear transformations

$$\varphi_t(x) = x + \sum_{l \geq 1} \beta_l(x) t^l$$

of $\mathfrak{g}$ and a formal (not necessarily invertible) parameter change $u = u(t)$ which transform the deformation $[x, y]_t$ into a deformation

$$[x, y]'_u = [x, y] + \sum_{k \geq 1} \alpha'_k(x, y) u^k, \quad \varphi_t^{-1}[\varphi_t(x), \varphi_t(y)]_t = [x, y]'_{u(t)}$$

with the cocycle $\alpha'_1 \in C^2(\mathfrak{g}, \mathfrak{g})$ being not cohomologous to 0. Otherwise the deformation is called singular.

We have already remarked that one can associate to an arbitrary $\mathbb{Z}_{>0}$ -deformation $\Psi = \Psi_1 + \Psi_2 + \Psi_3 + \dots$ of $\mathcal{V}_n$ the following one-parametric deformation:

$$[x, y]_t^\Psi = [x, y] + t\Psi_1(x, y) + t^2\Psi_2(x, y) + \dots + t^k\Psi_k(x, y) + \dots$$

One can also remark that in our case a formal deformation $[x, y]_t^\Psi$ is in fact a polynomial on t .

Proposition 8.4. *Let fix the isomorphism $O(UT_n, V_n) = \mathbb{K}^5$, $n \geq 16$, then a formal deformation $[x, y]_t^\Psi$ corresponding to some $\mathbb{Z}_{>0}$ -deformation Ψ of $\mathcal{V}_n$ is non-singular if and only if its orbit O_Ψ belongs to the union $\bigcup_{i=1}^5 Ox_i$ of coordinate lines Ox_i in $\mathbb{K}^5$.*

Acting by corresponding $\varphi_t = id + t\varphi_1 + t^2\varphi_2 + \dots$ we reduce our problem to a parameter change in

$$\Psi_t = t^{n-11}x_1\psi_{n,11} + t^{n-10}x_2\psi_{n,10} + \dots + t^{n-7}x_5\psi_{n,7}.$$

If $x_1 \neq 0$ then a parameter change $u(t) = t^{n-11}$ is possible iff $x_2 = x_3 = x_4 = x_5 = 0$ the other cases are treated in the same way.

Remark. If $n \geq 14$ one can associate to an arbitrary deformation of the form $\Psi = \sum_{i=1}^5 x_i\psi_{n,12-i}$ a linear one-parameter deformation $[\cdot]_\tau'$ of $\mathcal{V}_n$:

$$[x, y]'_\tau = [x, y] + \tau\Psi(x, y).$$

It follows from $[\Psi, \Psi] = 0$ in $C^3(\mathcal{V}_n, \mathcal{V}_n)$. Evidently a linear one-parameter deformation $[\cdot]_\tau'$ is non-singular iff $\Psi \neq 0$.

9. SYMPLECTIC STRUCTURES AND ONE-DIMENSIONAL CENTRAL EXTENSIONS

First of all we are going to calculate $H^2(\mathfrak{g})$ of an arbitrary $\mathbb{Z}_{>0}$ -deformation $\mathfrak{g} = (\mathcal{V}_n, [\cdot, \cdot] + \Psi)$. As we remarked in the section 4 the canonical basis $e_1, e_2, \dots, e_n$ of $\mathfrak{g} = (\mathcal{V}_n, [\cdot, \cdot] + \Psi)$ defines the filtration L of $\mathfrak{g}$:

$$\mathfrak{g} = L^1\mathfrak{g} \supset L^2\mathfrak{g} = \text{Span}(e_2, \dots, e_n) \supset \dots \supset L^n\mathfrak{g} = \text{Span}(e_n) \supset \{0\},$$

and L defines the filtration $\tilde{L}$ of the cochain complex $(C^*(\mathfrak{g}), d)$. For the simplicity one can assume that $\Psi = x_1\psi_{n,11} + \dots + x_5\psi_{n,7}$. Let us consider the corresponding to $\tilde{L}$ spectral sequence $E_r^{p,q}$ that converges to $H^*(\mathfrak{g})$. We recall also that

$$E_1^{p,q} = H_{(-p)}^{p+q}(\text{gr}_L\mathfrak{g}) = H_{(-p)}^{p+q}(\mathcal{V}_n).$$

We recall that the space $H^2(\mathcal{V}_n)$ is 3-dimensional and it is spanned by classes

$$g_5 = [e^2 \wedge e^3], \quad g_7 = [e^2 \wedge e^5 - 3e^3 \wedge e^4], \quad [\Omega_{n+1}] = \frac{1}{2} \left[\sum_{i+j=n+1} (j-i)e^i \wedge e^j \right]$$

of weights 5, 7, $n+1$ respectively. It is evident that the classes $[g_5]$ and $[g_7]$ survive to the term E_∞ because $d_{\tilde{\Psi}} e^i = de^i, i = 1, \dots, 7, \forall \tilde{\Psi}$. And the question is: does $[\Omega_{n+1}] \in E_1^{-2k-1, 2k+3}$ survive to E_∞ ?

Proposition 9.1. *Let $\mathfrak{g} = (\mathcal{V}_n, [,] + \tilde{\Psi})$ be a Lie algebra defined by the following deformation of $\mathcal{V}_n$:*

$$\tilde{\Psi} = \tilde{\Psi}_{n-11} + \dots + \tilde{\Psi}_{n-7} = x_1 \psi_{n,11} + \dots + x_5 \psi_{n,7}.$$

Then in the spectral sequence $E_r^{p,q}$ the following properties hold on:

- 1) $d_r \equiv 0, r = 1, \dots, n-12$;
- 2) $E_r^{p,q} = E_1^{p,q}, r = 2, \dots, n-11$;
- 3) $d_{n-11}(\Omega_{n+1}) = 0$ if and only if $x_1 = 0$.
- 4) If $x_1 = 0$ then $d_r(\Omega_{n+1}) = 0 \forall r \geq 1$, i.e the class $[\Omega_{n+1}]$ survive to the term E_∞ .

The differential d_Ψ of the cochain complex $(C^*(\mathfrak{g}^*), d)$ has the form:

$$d = d_0 + \Psi_{n-11}^* + \dots + \Psi_{n-7}^* = d_0 + x_1 \psi_{n,11}^* + \dots + x_5 \psi_{n,7}^*,$$

where by Ψ_i^* and $\psi_{n,j}^*$ we denote the dual applications to Ψ_i and $\psi_{n,j}$ respectively and d_0 stands for the differential of $(C^*(\mathcal{V}_n), d_0)$. The items 1) and 2) of the proposition are evident and

$$d_{n-11}([\Omega_{n+1}]) = [\Psi_{n-11}^*(\Omega_{n+1})] = x_1 [\psi_{n,11}^*(\Omega_{n+1})],$$

On another hand one can compute $\psi_{n,11}^*(\Omega_{n+1})$ explicitly:

$$(40) \quad \begin{aligned} -\psi_{n,11}^*(\Omega_{n+1}) = & -\psi_{n,11}^* \left((n-1)e^1 \wedge e^n + (n-3)e^2 \wedge e^{n-1} + \dots + (n-9)e^5 \wedge e^{n-4} \right) = \\ & (n-1)e^1 \wedge (Z_1(n-4)e^2 \wedge e^9 + Z_2(n-4)e^3 \wedge e^8 + Z_3(n-4)e^4 \wedge e^7 + Z_4(n-4)e^5 \wedge e^6) + \\ & (n-3)e^2 \wedge (Q_1(n-4)e^2 \wedge e^8 + Q_2(n-4)e^3 \wedge e^7 + Q_3(n-4)e^4 \wedge e^6) + \\ & + (n-5)e^3 \wedge (P_1(n-4)e^2 \wedge e^7 + P_2(n-4)e^3 \wedge e^6 + P_3(n-4)e^4 \wedge e^5) + \\ & + (n-7)e^4 \wedge \frac{n-5}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5) + (n-9)e^5 \wedge (e^2 \wedge e^5 - 3e^3 \wedge e^4) \end{aligned}$$

As it was shown in the Section 6 this element is cohomologous to

$$-\frac{1}{5544}(n-12)(n^2-12n+59)(n^2-21n+116)[2e^2 \wedge e^3 \wedge e^7 - 5e^2 \wedge e^4 \wedge e^6 + 20e^3 \wedge e^4 \wedge e^5]$$

and it defines a non-trivial element in $H_{(12)}^3(\mathcal{V}_n)$ for $n > 12$.

We can finish the proof by remark that

$$E_r^{-n-1+r, n-r+4} = E_1^{-n-1+r, n-r+4} = H_{(n+1-r)}^3(\mathcal{V}_n) = 0, r > n-11,$$

because $H_{(i)}^3(\mathcal{V}_n) = 0$ for all $i < 12$. Hence all other differentials of the spectral sequence

$$d_r : E_r^{-n-1, n+3} \rightarrow E_r^{-n-1+r, n-r+4}, r > n-11$$

are trivial.

Corollary 9.2. *Let $\mathfrak{g} = (\mathcal{V}_{2k}, [,] + \Psi)$ be a $\mathbb{Z}_{>0}$ -deformation of $\mathcal{V}_{2k}$ and $(x_1, x_2, \dots, x_5)$ be the set of homogeneous coordinates of the UT_n -orbit O_Ψ . $\mathfrak{g}$ admits a symplectic structure if and only if $x_1 = 0$.*

Corollary 9.3. *If the orbit O_Ψ of a $\mathbb{Z}_{>0}$ -deformation $\mathfrak{g} = (\mathcal{V}_n, [,] + \Psi$ has a non-zero first coordinate $x_1 \neq 0$ then:*

$$\dim H^1(\mathfrak{g}) = \dim H^2(\mathfrak{g}) = 2.$$

Hence for a generic deformations $\mathfrak{g}$ of $\mathcal{V}_n$ the Dixmier inequalities (see [5]) $\dim H^i(\mathfrak{g}) \geq 2$ are sharp for $i = 1, 2$.

Recall that a one-dimensional central extension of a Lie algebra $\mathfrak{g}$ is an exact sequence

$$(41) \quad 0 \rightarrow \mathbb{K} \rightarrow \tilde{\mathfrak{g}} \rightarrow \mathfrak{g} \rightarrow 0$$

of Lie algebras and their homomorphisms, in which the image of the homomorphism $\mathbb{K} \rightarrow \tilde{\mathfrak{g}}$ is contained in the center of the Lie algebra $\mathfrak{g}$. To the cocycle $c \in \Lambda^2(\mathfrak{g}^*)$ corresponds the extension

$$0 \rightarrow \mathbb{K} \rightarrow \mathbb{K} \oplus \mathfrak{g} \rightarrow \mathfrak{g} \rightarrow 0$$

where the Lie bracket in $\mathbb{K} \oplus \mathfrak{g}$ is defined by the formula

$$[(\lambda, g), (\mu, h)] = (c(g, h), [g, h]).$$

It can be checked directly that the Jacobi identity for this Lie bracket is equivalent to c being cocycle and that to cohomologous cocycles correspond equivalent (in a obvious sense) extensions.

Now let $\tilde{\mathfrak{g}}$ be a filiform Lie algebra, it has one-dimensional center $Z(\tilde{\mathfrak{g}})$ and we have the following one-dimensional central extension:

$$0 \rightarrow \mathbb{K} = Z(\tilde{\mathfrak{g}}) \rightarrow \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}}/Z(\tilde{\mathfrak{g}}) \rightarrow 0$$

where the quotient Lie algebra $\tilde{\mathfrak{g}}/Z(\tilde{\mathfrak{g}})$ is also filiform Lie algebra. Moreover, let $e_1, \dots, e_{n+1}$ be some adapted basis of $\tilde{\mathfrak{g}}$ ($Z(\tilde{\mathfrak{g}}) = \text{Span}(e_{n+1})$) and $e^1, \dots, e^{n+1} \in \mathfrak{g}^*$ its dual basis. Then the forms $e^1, \dots, e^n$ can be regarded as the dual basis to the basis $e_1 + Z(\tilde{\mathfrak{g}}), \dots, e_n + Z(\tilde{\mathfrak{g}})$ of $\tilde{\mathfrak{g}}/Z(\tilde{\mathfrak{g}})$ and the cocycle $\Omega_{n+1} = de^{n+1}$ determines this one-dimensional central extension.

Proposition 9.4. *Let $\mathfrak{g}$ be a filiform Lie algebra and let its center $Z(\mathfrak{g})$ be spanned by some $\xi \in Z(\mathfrak{g})$. Then $\tilde{\mathfrak{g}}$ taken from a one-dimensional central extension*

$$0 \rightarrow \mathbb{K} \rightarrow \tilde{\mathfrak{g}} \rightarrow \mathfrak{g} \rightarrow 0$$

with cocycle $c \in \Lambda^2(\mathfrak{g})$ is a filiform Lie algebra if and only if the restricted function $f(\cdot) = c(\cdot, \xi)$ is non-trivial in $\mathfrak{g}^$.*

Corollary 9.5. *Let $\mathfrak{g}$ be a symplectic filiform Lie algebra and ω its symplectic form, then the one-dimensional central extension $\tilde{\mathfrak{g}}$ with the cocycle $c = \omega$ will be also a filiform Lie algebra.*

Theorem 9.6. *Let $\tilde{\mathfrak{g}} = (\mathcal{V}_{2k+1}, [,] + \tilde{\Psi})$ be $\mathbb{Z}_{>0}$ -filtered deformation of $\mathcal{V}_{2k+1}$, where*

$$\tilde{\Psi} = x_1\psi_{2k+1,11} + \dots + x_5\psi_{2k+1,7}.$$

Then $\tilde{\mathfrak{g}}$ can be represented as a one-dimensional central extension of $\mathbb{Z}_{>0}$ -filtered deformation $\mathfrak{g}_X = (\mathcal{V}_{2k}, [,] + \Phi)$ of $\mathcal{V}_{2k}$, where

$$\Phi = x_1\psi_{2k,10} + x_2\psi_{2k,9} + x_3\psi_{2k,8} + x_4\psi_{2k,7}, \quad X = (x_1, x - 2, x_3, x_4).$$

The cocycle Ω_{X,x_5} that determines this one-dimensional central extension is equal to

$$(42) \quad \begin{aligned} \Omega_{X,x_5} &= \frac{1}{2} \sum_{i+j=2k+1} (j-i)e^i \wedge e^j + x_1 \Omega_{2k,11} + x_2 \Omega_{2k,10} + x_3 \Omega_{2k,9} + x_4 \Omega_{2k,8} + x_5 \Omega_{2k,7}, \\ \Omega_{2k,11} &= Z_1(2k-3)e^2 \wedge e^9 + Z_2(2k-3)e^3 \wedge e^8 + Z_3(2k-3)e^4 \wedge e^7 + Z_4(2k-3)e^5 \wedge e^6, \\ \Omega_{2k,10} &= Q_1(2k-2)e^2 \wedge e^8 + Q_2(2k-2)e^3 \wedge e^7 + Q_3(2k-2)e^4 \wedge e^6, \\ \Omega_{2k,9} &= P_1(2k-1)e^2 \wedge e^7 + P_2(2k-1)e^3 \wedge e^6 + P_3(2k-1)e^4 \wedge e^5, \\ \Omega_{2k,8} &= \frac{2k-1}{2}(e^2 \wedge e^6 - 2e^3 \wedge e^5), \quad \Omega_{2k,7} = e^2 \wedge e^5 - 3e^3 \wedge e^4. \end{aligned}$$

Definition 9.7. Let $\mathfrak{g}, \tilde{\mathfrak{g}}$ be two symplectic Lie algebras, $\omega_{\mathfrak{g}}, \omega_{\tilde{\mathfrak{g}}}$ be corresponding symplectic structures. A Lie algebras isomorphism $f : \mathfrak{g} \rightarrow \tilde{\mathfrak{g}}$ is called a symplecto-isomorphism $f : (\mathfrak{g}, \omega_{\mathfrak{g}}) \rightarrow (\tilde{\mathfrak{g}}, \omega_{\tilde{\mathfrak{g}}})$ if and only if $\omega_{\mathfrak{g}} = f^*(\omega_{\tilde{\mathfrak{g}}})$.

Theorem 9.8. 1) Let $\mathfrak{g}$ be a symplectic filtered deformation of $\mathcal{V}_{2k}$, $\omega_{\mathfrak{g}}$ its symplectic structure. Then there exists a vector $X = (x_1, x_2, x_3, x_4) \in \mathbb{K}^4$, $x_5 \in \mathbb{K}$ such that the pair $(\mathfrak{g}, \omega_{\mathfrak{g}})$ is symplecto-isomorphic to $(\mathfrak{g}_X, \Omega_{X,x_5})$, where $\mathfrak{g}_X$ is one of the Lie algebras from the table below

$i+j$	$\mathfrak{g}_X$: commuting relations $[e_i, e_j]$, $i < j$
$3 \leq i+j \leq 6$	$[e_i, e_j] = (j-i)e_{i+j}$
7	$[e_1, e_6] = 5e_7,$ $[e_2, e_5] = 3e_7 + x_1 e_{2k-3} + x_2 e_{2k-2} + x_3 e_{2k-1} + x_4 e_n,$ $[e_3, e_4] = e_7 - 3x_1 e_{2k-3} - 3x_2 e_{2k-2} - 3x_3 e_{2k-1} - 3x_4 e_n,$
8	$[e_1, e_7] = 6e_8,$ $[e_2, e_6] = 4e_8 + x_1 \frac{(2k-4)}{2} e_{2k-2} + x_2 \frac{(2k-3)}{2} e_{2k-1} + x_3 \frac{(2k-2)}{2} e_{2k},$ $[e_3, e_5] = 2e_8 + x_1(2k-4)e_{2k-2} + x_2(2k-3)e_{2k-1} + x_3(2k-2)e_{2k};$
9	$[e_1, e_8] = 7e_9,$ $[e_2, e_7] = 5e_9 + x_1 \frac{(5\binom{2k-3}{2} + 3\binom{2k-3}{1} - 6)}{21} e_{2k-1} + x_2 \frac{(5\binom{2k-2}{2} + 3\binom{2k-2}{1} - 6)}{21} e_{2k},$ $[e_3, e_6] = 3e_9 - x_1 \frac{(4\binom{2k-3}{2} + 15\binom{2k-3}{1} - 30)}{21} e_{2k-1} - x_2 \frac{(4\binom{2k-2}{2} + 15\binom{2k-2}{1} - 30)}{21} e_{2k},$ $[e_4, e_5] = e_9 - x_1 \frac{(13\binom{2k-3}{2} - 30\binom{2k-3}{1} + 60)}{21} e_{2k-1} - x_2 \frac{(13\binom{2k-2}{2} - 30\binom{2k-2}{1} + 60)}{21} e_{2k-1},$
10	$[e_1, e_9] = 8e_{10},$ $[e_2, e_8] = 6e_{10} + x_1 \frac{(3\binom{2k-2}{3} + 4\binom{2k-2}{2} - 4\binom{2k-2}{1})}{28} e_{2k},$ $[e_3, e_7] = 4e_{10} + x_1 \frac{((\binom{2k-2}{3} - 8\binom{2k-2}{2}) + 8\binom{n-2}{1})}{14} e_{2k},$ $[e_4, e_6] = 2e_{10} + x_1 \frac{(-13\binom{2k-2}{3} + 20\binom{2k-2}{2} - 20\binom{2k-2}{1})}{28} e_n,$
$11 \leq i+j \leq 2k$	$[e_i, e_j] = (j-i)e_{i+j}$

and its corresponding symplectic form Ω_{X,x_5} is equal to

$$\Omega_{X,x_5} = \frac{1}{2} \sum_{i+j=2k+1} (j-i)e^i \wedge e^j + x_1 \Omega_{2k,11} + x_2 \Omega_{2k,10} + x_3 \Omega_{2k,9} + x_4 \Omega_{2k,8} + x_5 \Omega_{2k,7}.$$

2) A pair $(\mathfrak{g}_X, \Omega_{X, x_5})$ is symplecto-isomorphic to $(\mathfrak{g}_Y, \Omega_{Y, y_5})$ if and only if there exist an $\alpha \in \mathbb{K}^*$ such that

$$y_1 = \alpha^{n-11}x_1, y_2 = \alpha^{n-10}x_2, y_3 = \alpha^{n-9}x_3, y_4 = \alpha^{n-8}x_4, y_5 = \alpha^{n-7}x_5.$$

Remark. The previous theorem shows that all $\mathbb{Z}_{>0}$ -deformations $\tilde{\mathfrak{g}}$ of $\mathcal{V}_{2k+1}$ are contact Lie algebras and gives their complete classification. An arbitrary symplectic $\mathbb{Z}_{>0}$ -deformations $\mathfrak{g}$ of $\mathcal{V}_{2k}$ can be obtained as quotient Lie algebra $\tilde{\mathfrak{g}}/Z(\tilde{\mathfrak{g}})$. This method of classification of symplecto-isomorphism classes of low-dimensional filiform Lie algebras was considered in [10].

Taking rational coordinates $(x_1, x_2, \dots, x_5)$ in the table above one will get a nilpotent Lie algebra $\mathfrak{g}_X$ with rational structure constants and hence due to the Malcev theorem ([14]) the corresponding simply connected nilpotent Lie group G has a cocompact lattice Γ . Thus one will get a family of examples of symplectic nilmanifolds $M = G/\Gamma$.

REFERENCES

- [1] I. K. Babenko and I. A. Taimanov, *On the existence of nonformal simply connected symplectic manifolds*, Russian Math. Surveys, **53**:4 (1998), 1082–1083.
- [2] I. K. Babenko and I. A. Taimanov, *On nonformal simply connected symplectic manifolds*, Siberian Math. J., **41**:2 (2000), 204–217.
- [3] V. M. Bukhshtaber, *Groups of polynomial transformations of a line, nonformal symplectic manifolds, and the landweber-Novikov algebra*, Russian Math. Surveys, **54**:4 (1999), 837–838.
- [4] B. Chu, *Symplectic homogeneous spaces*, Trans. Amer. Math. Soc., **197** (1974), 145–159.
- [5] J. Dixmier *Cohomologie des algèbres de Lie nilpotentes*, Acta sci. math. Szeged **16** (1955), 246–250.
- [6] B. Feigin, D. Fuchs, *Homology of the Lie algebras of vector fields on the line*, Funct. Anal. Appl., **14**:3 (1980), 45–60.
- [7] A. Fialowski, *Deformations of the Lie algebra of vector fields on the line*, Russian Math. Surveys, **38**:1 (1983), 185–186.
- [8] A. Fialowski, D. Fuchs, *Singular deformations of Lie algebras. Example: deformations of the Lie algebra L_1* , in: Topics in singularity theory (V.I. Arnold 60th anniversary collection), Amer. Math. Soc. Transl. Ser.2, **180** (1997), Amer. Math. Soc., Providence, RI, 77–92.
- [9] D. Fuchs, *Cohomology of infinite-dimensional Lie algebras* Consultants Bureau, N.Y., London, 1986.
- [10] J.R. Gómez, A. Jimenéz-Merchán, Y. Khakimdjano, *Symplectic structures on the filiform Lie algebras*, J.Pure Appl.Algebra, **156** (2001), 15–31.
- [11] M. Goze, A. Bouyakoub, *Sur les algèbres de Lie munies d’une forme symplectique*, Rend. Sem. Fac. Sc. Univ. Cagliari, **57**:1 (1987), 85–97.
- [12] M. Goze, Y. Khakimdjano, *Nilpotent Lie algebras*, Kluwer Academic, Dordrecht, 1996.
- [13] Yu. Khakimdjano, *Varieties of Lie algebras laws*, in Handbook of algebra, vol.2, 509–541, North-Holland, Amsterdam, 2000.
- [14] A. Malcev, *On a class of homogeneous spaces*, Izvestia Akad. Nauk SSSR Ser. Mat. **3** (1949), 9–32 (Russian); English translation: Amer. Math. Soc. Transl. (1) **9** (1962), 276–307.
- [15] A. Médina, P. Revoy, *Groupes de Lie à structure de symplectique invariante*, in "Symplectic geometry, groupoids and integrable systems. Séminaire Sud Rhodanien." M.S.R.I., Springer-Verlag, New York / Berlin (1991), 247–266.
- [16] D.V. Millionschikov, *Cohomology of nilmanifolds and Gontcharova’s theorem*, Russian Math. Surveys, **56**:4 (2001), 758–759 (in russian).
- [17] D.V. Millionschikov, *Cohomology of nilmanifolds and Gontcharova’s theorem*, in "Global Differential geometry: The Mathematical Legacy of Alfred Gray", M. Fernandez and J.Wolf ed., AMS CONM 288 (2001), 381–385.
- [18] D.V. Millionschikov, *N-graded filiform Lie algebras*, Uspekhi Mat. Nauk, **57**:2 (2002), 422–424.
- [19] D.V. Millionschikov, *N-graded filiform Lie algebras and symplectic structures*, math.RA/0205042.

- [20] V. Morosov, *Classification of nilpotent Lie algebras of order 6*, Izv. Vyssh. Uchebn. Zaved. Mat., **4** (1958), 161–171.
- [21] A. Nijenhuis, R.W. Richardson, Jr., *Deformations of Lie algebra structures*, J. of Math. and Mech., **17**:1 (1967), 89–105.
- [22] M. Vergne, *Réductibilité de la variété des algèbres de Lie nilpotentes*, C.R. Acad. Sc. Paris. **263** (1966), 4–6.
- [23] M. Vergne, *Cohomologie des algèbres de Lie nilpotentes*, Bull. Soc. Math. France **98** (1970), 81–116.

DEPARTMENT OF MATHEMATICS AND MECHANICS, MOSCOW STATE UNIVERSITY, 119899 MOSCOW, RUSSIA

Current address: Université Louis Pasteur, UFR de Mathématique et d'Informatique, 7 rue René Descartes - 67084 Strasbourg Cedex (France)

E-mail address: `million@mech.math.msu.su`